



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07K 14/60, 5/02, 5/06, A61K 38/05, 38/25	A1	(11) International Publication Number: WO 00/01726 (43) International Publication Date: 13 January 2000 (13.01.00)
(21) International Application Number: PCT/DK99/00368 (22) International Filing Date: 29 June 1999 (29.06.99) (30) Priority Data: PA 1998 00857 30 June 1998 (30.06.98) DK PA 1998 01440 9 November 1998 (09.11.98) DK (71) Applicant: NOVO NORDISK A/S [DK/DK]; Novo Allé, DK-2880 Bagsvaerd (DK). (72) Inventors: PESCHKE, Bernd; Eskebjerggaard 56, DK-2760 Maaloev (DK). RICHTER, Stefan, Lutz (deceased). HANSEN, Thomas, Kruse; Tibbevangen 78, DK-2730 Herlev (DK). ANKERSEN, Michael; 1.th Dalgas Have 34, DK-2000 Frederiksberg (DK).		(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i>
(54) Title: COMPOUNDS WITH GROWTH HORMONE RELEASING PROPERTIES (57) Abstract <p>The invention relates to novel compounds, compositions containing them, and their use for treating medical disorders resulting from a deficiency in growth hormone.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon	KR	Republic of Korea	PL	Poland		
CN	China	KZ	Kazakhstan	PT	Portugal		
CU	Cuba	LC	Saint Lucia	RO	Romania		
CZ	Czech Republic	LI	Liechtenstein	RU	Russian Federation		
DE	Germany	LK	Sri Lanka	SD	Sudan		
DK	Denmark	LR	Liberia	SE	Sweden		
EE	Estonia			SG	Singapore		

COMPOUNDS WITH GROWTH HORMONE RELEASING PROPERTIES

FIELD OF INVENTION

5

The present invention relates to novel compounds, compositions containing them, and their use for treating medical disorders resulting from a deficiency in growth hormone.

BACKGROUND OF THE INVENTION

10

Growth hormone is a hormone which stimulates growth of all tissues capable of growing. In addition, growth hormone is known to have a number of effects on metabolic processes, e.g., stimulation of protein synthesis and free fatty acid mobilisation and to cause a switch in energy metabolism from carbohydrate to fatty acid metabolism. Deficiency in growth hormone can result in a number of severe medical disorders, e.g., dwarfism.

15

Growth hormone is released from the pituitary. The release is under tight control of a number of hormones and neurotransmitters either directly or indirectly. Growth hormone release can be stimulated by growth hormone releasing hormone (GHRH) and inhibited by somatostatin. In both cases the hormones are released from the hypothalamus but their action is mediated primarily via specific receptors located in the pituitary. Other compounds which stimulate the release of growth hormone from the pituitary have also been described. For example arginine, L-3,4-dihydroxyphenylalanine (L-Dopa), glucagon, vasopressin, PACAP (pituitary adenylyl cyclase activating peptide), muscarinic receptor agonists and a synthetic hexapeptide, GHRP (growth hormone releasing peptide) release endogenous growth hormone either by a direct effect on the pituitary or by affecting the release of GHRH and/or somatostatin from the hypothalamus.

20

25

In disorders or conditions where increased levels of growth hormone is desired, the protein nature of growth hormone makes anything but parenteral administration non-viable. Furthermore, other directly acting natural secretagogues, e.g., GHRH and PACAP, are longer polypeptides for which reason parenteral administration is preferred.

30

The use of certain compounds for increasing the levels of growth hormone in mammals has previously been proposed, e.g. in EP 18 072, EP 83 864, WO 8302272, WO 8907110, WO 8901711, WO 8910933, WO 8809780, WO 9118016, WO 9201711, WO 9304081, WO 9413696, WO 9517423, WO 9514666, WO 9615148, WO 9622997, WO 9635713, WO 9700894, WO 9722620, WO 9723508, WO 9740023, and WO 9810653.

The composition of growth hormone releasing compounds is important for their growth hormone releasing potency as well as their bioavailability. It is therefore an object of the present invention to provide novel compounds with growth hormone releasing properties. Moreover, it is an object to provide novel growth hormone releasing compounds (growth hormone secretagogues) which are specific and/or selective and have no or substantially no side-effects, such as e.g. release of LH, FSH, TSH, ACTH, vasopressin, oxytocin, cortisol and/or prolactin. It is also an object to provide compounds which have good oral bioavailability.

SUMMARY OF THE INVENTION


In accordance with the present invention there is provided novel compounds which act directly on the pituitary cells under normal experimental conditions in vitro to release growth hormone therefrom.

These growth hormone releasing compounds can be utilized in vitro as unique research tools for understanding, inter alia, how growth hormone secretion is regulated at the pituitary level.

Moreover, the growth hormone releasing compounds of the present invention can also be administered in vivo to increase endogenous growth hormone release.

DESCRIPTION OF THE INVENTION

Accordingly, the present invention relates to a compound of the general formula I




formula I

wherein


R¹ is hydrogen or C₁₋₆-alkyl;

R² is hydrogen or C₁₋₆-alkyl;

L is



or



wherein R⁴ is hydrogen or C₁₋₆ alkyl;

p is 0 or 1;

q, s, t, u are independently from each other 0, 1, 2, 3 or 4;


r is 0 or 1;

the sum $q + r + s + t + u$ is 0, 1, 2, 3, or 4;

R^9 , R^{10} , R^{11} , and R^{12} are independently from each other hydrogen or C_{1-6} alkyl;

5

Q is $>N-R^{13}$ or



wherein o is 0, 1 or 2;


10

T is $-N(R^{15})(R^{16})$ or hydroxyl;

R^{13} , R^{15} , and R^{16} are independently from each other hydrogen or C_{1-6} alkyl;


15 R^{14} is hydrogen, aryl or hetaryl;

G is $-O-(CH_2)_k-R^{17}$,



20 wherein R^{17} , R^{18} , R^{19} , R^{20} and R^{21} independently from each other are hydrogen, halogen, aryl, hetaryl, C_{1-6} -alkyl or C_{1-6} -alkoxy;
 k is 0, 1 or 2;

J is $-O-(CH_2)_I-R^{22}$,



5 wherein R^{22} , R^{23} , R^{24} , R^{25} and R^{26} independently from each other are hydrogen, halogen, aryl, hetaryl, C_{1-6} -alkyl or C_{1-6} -alkoxy;

I is 0, 1 or 2;

a is 0, 1, or 2;

10

b is 0, 1, or 2;

c is 0, 1, or 2;

15 d is 0 or 1;

e is 0, 1, 2, or 3;

f is 0 or 1;

20

R^5 is hydrogen or C_{1-6} -alkyl optionally substituted with one or more hydroxyl, aryl or hetaryl;

R^6 and R^7 are independently from each other hydrogen or C_{1-6} -alkyl, optionally substituted with one or more halogen, amino, hydroxyl, aryl, or hetaryl;

25

R^8 is hydrogen or C_{1-6} -alkyl, optionally substituted with one or more halogen, amino, hydroxyl, aryl, or hetaryl;

R^6 and R^7 or R^6 and R^8 or R^7 and R^8 may optionally form $-(CH_2)_i-U-(CH_2)_j-$, wherein i and j independently from each other are 1, 2 or 3 and U is $-O-$, $-S-$, or a valence bond;

5 M is arylene, hetarylene, $-O-$, $-S-$ or $-CR^{27}=CR^{28}-$;

R^{27} and R^{28} are independently from each other hydrogen or C_{1-6} -alkyl, optionally substituted with one or more aryl or hetaryl;

or a pharmaceutically acceptable salt thereof.

10

Moreover, the compounds of formula I may comprise any optical isomers thereof, in the form of separated, pure or partially purified optical isomers or racemic mixtures thereof. Whenever one or more chiral carbon atoms are present such chiral center or centers may be in the R- and/or S-configuration, or a mixture of R and S.

15


Furthermore, the compounds of formula I may have one or more carbon-carbon double bonds with the possibility of geometric isomeri, and it is intended that possible stereoisomers (E or Z isomers) are included in the scope of the invention, unless a special geometric isomer is specified.

20

In one embodiment of the compound of formula I R^1 is C_{1-6} -alkyl, such as C_{1-4} -alkyl, in particular methyl. In a second embodiment R^1 is hydrogen.

25 In a further embodiment of the compound of formula I R^2 is C_{1-6} -alkyl, such as C_{1-4} -alkyl, in particular methyl.

In a still further embodiment of the compound of formula I L is



wherein R^4 is hydrogen or C_{1-6} alkyl;

p is 0 or 1;


q, s, t, u are independently from each other 0, 1, 2, 3 or 4;

5 r is 0 or 1;

the sum $q + r + s + t + u$ is 0, 1, 2, 3, or 4;

R^9 , R^{10} , R^{11} , and R^{12} are independently from each other hydrogen or C_{1-6} alkyl;

Q is $>N-R^{13}$ or



10 wherein o is 0, 1 or 2;

T is $-N(R^{15})(R^{16})$ or hydroxyl;

R^{13} , R^{15} , and R^{16} are independently from each other hydrogen or C_{1-6} alkyl;

R^{14} is hydrogen, aryl or hetaryl. In one embodiment R^4 is hydrogen. In a second embodiment

R^4 is C_{1-6} -alkyl, such as C_{1-4} -alkyl, in particular methyl. In a third embodiment p is 0. In a fur-

15 ther embodiment q is 0. In a still further embodiment q is 1. In a further embodiment s is 0. In a still further embodiment s is 1. In a further embodiment t is 0. In a still further embodiment t

is 1. In a further embodiment u is 0. In a still further embodiment u is 1. In a further embodi-

ment r is 0. In a still further embodiment r is 1. In a further embodiment R^9 is hydrogen. In a

20 still further embodiment R^9 is C_{1-6} -alkyl, such as C_{1-4} -alkyl, in particular methyl. In a further

embodiment R^{10} is hydrogen. In a still further embodiment R^{10} is C_{1-6} -alkyl, such as C_{1-4} -alkyl,


in particular methyl. In a further embodiment R^{11} is hydrogen. In a still further embodiment

R^{11} is C_{1-6} -alkyl, such as C_{1-4} -alkyl, in particular methyl. In a further embodiment R^{12} is hydro-

gen. In a still further embodiment R^{12} is C_{1-6} -alkyl, such as C_{1-4} -alkyl, in particular methyl. In a

further embodiment Q is $>N-R^{13}$. In a still further embodiment R^{13} is hydrogen. In a further


25 embodiment R^{13} is C_{1-6} -alkyl, such as C_{1-4} -alkyl, in particular methyl. In a still further embodi-



In a still further embodiment R^{14} is hetaryl, in particular thiazolyl. In a further embodiment R^{14} is hydrogen. In a still further embodiment o is 0. In a further embodiment o is 1. In a still fur-

ther embodiment T is hydroxyl. In a further embodiment T is $-N(R^{15})(R^{16})$. In a still further embodiment R^{15} is C_{1-6} -alkyl, such as C_{1-4} -alkyl, in particular methyl. In a further embodiment R^{16} is C_{1-6} -alkyl, such as C_{1-4} -alkyl, in particular methyl.

- 5 In a further embodiment of the compound of formula I L is




wherein q, s, t, u independently from each other are 0, 1, 2, 3 or 4;

r is 0 or 1;

- 10 the sum $q + r + s + t + u$ is 0, 1, 2, 3, or 4;

R^9 , R^{10} , R^{11} , and R^{12} are independently from each other hydrogen or C_{1-6} alkyl;

Q is $>N-R^{13}$ or



wherein o is 0, 1 or 2;


- 15 T is $-N(R^{15})(R^{16})$ or hydroxyl;

R^{13} , R^{15} , and R^{16} are independently from each other hydrogen or C_{1-6} alkyl;

R^{14} is hydrogen, aryl or hetaryl. In one embodiment q is 0. In a second embodiment q is 1. In a third embodiment s is 0. In a further embodiment s is 1. In a still further embodiment t is 0. In a further embodiment t is 1. In a still further embodiment u is 0. In a further embodiment u

- 20 is 1. In a still further embodiment r is 0. In a further embodiment r is 1. In a still further embodiment R^9 is hydrogen. In a still further embodiment R^9 is C_{1-6} -alkyl, such as C_{1-4} -alkyl, in particular methyl. In a further embodiment R^{10} is hydrogen. In a still further embodiment R^{10} is C_{1-6} -alkyl, such as C_{1-4} -alkyl, in particular methyl. In a further embodiment R^{11} is hydrogen. In a still further embodiment R^{11} is C_{1-6} -alkyl, such as C_{1-4} -alkyl, in particular methyl. In a further embodiment R^{12} is hydrogen. In a still further embodiment R^{12} is C_{1-6} -alkyl, such as C_{1-4} -
- 25

alkyl, in particular methyl. In a further embodiment Q is $>N-R^{13}$. In a still further embodiment R^{13} is hydrogen. In a still further embodiment R^{13} is C_{1-6} -alkyl, such as C_{1-4} -alkyl, in particular methyl. In a further embodiment Q is




- 5 In a still further embodiment R^{14} is hetaryl, in particular thiazolyl. In a further embodiment R^{14} is hydrogen. In a still further embodiment o is 0. In a further embodiment o is 1. In a still further embodiment T is hydroxyl. In a further embodiment T is $-N(R^{15})(R^{16})$. In a still further embodiment R^{15} is C_{1-6} -alkyl, such as C_{1-4} -alkyl, in particular methyl. In a further embodiment R^{16} is C_{1-6} -alkyl, such as C_{1-4} -alkyl, in particular methyl.

10

In the compound of the above formula I L is preferably 4-hydroxy-4-(2-thienyl)piperidino, (3-hydroxycyclohexyl)amino, 4-(N,N-dimethylamino)piperidino, N-methyl-N-(1-methylpiperidin-4-yl)amino, 4-((N,N-dimethylamino)methyl)piperidino, 4-methylpiperazino, (2,2,6,6-tetramethylpiperidine-4-yl)amino, 4-hydroxypiperidino, (3S)-3-((N,N-dimethylamino)methyl)-piperidino, (2S)-2-((N,N-dimethylamino)methyl)pyrrolidino,

15


In a still further embodiment of the compound of formula I G is



- 20 wherein R^{17} , R^{18} , R^{19} , R^{20} and R^{21} independently from each other are hydrogen, halogen, aryl, hetaryl, C_{1-6} -alkyl or C_{1-6} -alkoxy. In one embodiment R^{17} is hydrogen. In a second embodiment R^{18} is hydrogen. In a third embodiment R^{19} is hydrogen. In a further embodiment R^{19} is aryl, in particular phenyl. In a still further embodiment R^{20} is hydrogen. In a further embodiment R^{21} is hydrogen. In the compound of the above formula I G is preferably 2-naphthyl or biphenyl-4-yl.

25

In a further embodiment of the compound of formula I J is



- 5 wherein R^{22} , R^{23} , R^{24} , R^{25} and R^{26} independently from each other are hydrogen, halogen, aryl, hetaryl, C_{1-6} -alkyl or C_{1-6} -alkoxy. In one embodiment R^{22} is hydrogen. In a second embodiment R^{23} is hydrogen. In a third embodiment R^{24} is hydrogen. In a further embodiment R^{24} is halogen, in particular fluorine. In a still further embodiment R^{25} is hydrogen. In a further embodiment R^{26} is hydrogen. In the compound of the above formula I J is preferably phenyl, 4-fluoro-phenyl or 2-thienyl.
- 10

In a still further embodiment of the compound of formula I a is 1.

In a further embodiment of the compound of formula I b is 1.

15

In a still further embodiment of the compound of formula I c is 0.

In a further embodiment of the compound of formula I d is 0.

- 20 In a still further embodiment of the compound of formula I M is arylene or $-CR^{27}=CR^{28}-$, wherein R^{27} and R^{28} independently from each other are hydrogen or C_{1-6} -alkyl, optionally substituted with aryl or hetaryl. In one embodiment M is arylene, in particular phenylene. In another embodiment M is $-CR^{27}=CR^{28}-$, wherein R^{27} and R^{28} are independently hydrogen or C_{1-6} -alkyl. In a further embodiment R^{27} is hydrogen. In a still further embodiment
- 25 R^{27} is C_{1-6} -alkyl, in particular methyl. In a further embodiment R^{28} is hydrogen. In a further embodiment M is the E-isomer of $-CR^{27}=CR^{28}-$. In the compound of the above formula I M is preferably ethenylene, 1,3-phenylene or 1,2-propenylene.

In a further embodiment of the compound of formula I e is 0.

In a still further embodiment of the compound of formula I e is 1.

In a further embodiment of the compound of formula I f is 0.

5

In a still further embodiment of the compound of formula I f is 1.

In a further embodiment of the compound of formula I R^6 and R^7 are independently from each other hydrogen or C_{1-6} -alkyl. In one embodiment R^6 is hydrogen. In a second
10 embodiment R^6 is C_{1-6} -alkyl, in particular methyl. In a third embodiment R^7 is hydrogen. In a further embodiment R^7 is C_{1-6} -alkyl, in particular methyl.


In a still further embodiment of the compound of formula I R^6 and R^7 or R^6 and R^8 or R^7 and R^8 may optionally form $-(CH_2)_i-U-(CH_2)_j-$, wherein i and j independently from each other are 1 or
15 2 and U is -O-, -S-, or a valence bond.

In a further embodiment of the compound of formula I R^6 and R^7 form $-(CH_2)_i-U-(CH_2)_j-$, wherein i and j independently from each other are 1, 2 or 3 and U is -O-, -S-, or a valence bond. In one embodiment the sum i + j is 3. In a second embodiment U is a valence bond. In
20 a particular embodiment (CR^6R^7) is cyclobutyl.

In a still further embodiment of the compound of formula I R^6 and R^7 form $-(CH_2)_i-U-(CH_2)_j-$, wherein i and j independently from each other are 1 or 2 and U is -O-, -S-, or a valence bond. In one embodiment the sum i + j is 3. In a second embodiment U is a valence bond. In
25 a particular embodiment (CR^6R^7) is cyclobutyl.

In a further embodiment of the compound of formula I R^8 is hydrogen. In a second embodiment R^8 is C_{1-6} -alkyl, in particular methyl.

30 In a special embodiment the present invention relates to a compound of the general formula I



formula I


wherein

5


R^1 is hydrogen or C_{1-6} -alkyl;

R^2 is C_{1-6} -alkyl;

10 L is



or



wherein R^4 is hydrogen or C_{1-6} alkyl;

15 p is 0 or 1;

q, s, t, u are independently from each other 0, 1, 2, 3 or 4;


r is 0 or 1;

20

the sum $q + r + s + t + u$ is 0, 1, 2, 3, or 4;

R^9 , R^{10} , R^{11} , and R^{12} are independently from each other hydrogen or C_{1-6} alkyl;

Q is $>\text{N-R}^{13}$ or



wherein o is 0, 1 or 2;


5

T is $-\text{N}(\text{R}^{15})(\text{R}^{16})$ or hydroxyl;

R^{13} , R^{15} , and R^{16} are independently from each other hydrogen or C_{1-6} alkyl;


10 R^{14} is hydrogen, aryl or hetaryl;

G is



15 wherein R^{17} , R^{18} , R^{19} , R^{20} and R^{21} independently from each other are hydrogen, halogen, aryl, hetaryl, C_{1-6} -alkyl or C_{1-6} -alkoxy;

J is



20

wherein R^{22} , R^{23} , R^{24} , R^{25} and R^{26} independently from each other are hydrogen, halogen, aryl, hetaryl, C_{1-6} -alkyl or C_{1-6} -alkoxy;

a is 0, 1, or 2;

b is 0, 1, or 2;

5 c is 0, 1, or 2;

d is 0 or 1;

e is 0, 1, 2, or 3;

10

f is 0 or 1;

R⁵ is hydrogen or C₁₋₆-alkyl optionally substituted with one or more hydroxyl, aryl or hetaryl;

15 R⁶ and R⁷ are independently from each other hydrogen or C₁₋₆-alkyl, optionally substituted with one or more halogen, amino, hydroxyl, aryl, or hetaryl;

R⁸ is hydrogen or C₁₋₆-alkyl, optionally substituted with one or more halogen, amino, hydroxyl, aryl, or hetaryl;

20

R⁶ and R⁷ or R⁶ and R⁸ or R⁷ and R⁸ may optionally form -(CH₂)_i-U-(CH₂)_j-, wherein i and j independently from each other are 1, 2 or 3 and U is -O-, -S-, or a valence bond;

M is arylene or -CR²⁷=CR²⁸-;


25

R²⁷ and R²⁸ are independently from each other hydrogen or C₁₋₆-alkyl, optionally substituted with one or more aryl or hetaryl;
or a pharmaceutically acceptable salt thereof.


30 Preferred compounds of formula I of the invention are:

(2E)-5-Amino-5-methylhex-2-enoic acid N-((1R)-1-{N-[(1R)-1-benzyl-2-(4-((dimethylamino)-methyl)piperidin-1-yl)-2-oxoethyl]-N-methylcarbamoyl}-2-(2-naphthyl)ethyl)-N-methylamide


15



(2E)-5-Amino-5-methylhex-2-enoic acid N-((1R)-1-{N-[(1R)-1-benzyl-2-((3S)-3-(dimethylaminomethyl)piperidin-1-yl)-2-oxoethyl]-N-methylcarbamoyl}-2-(2-naphthyl)ethyl)-N-methylamide




(2E)-4-(1-Aminocyclobutyl)but-2-enoic acid N-((1R)-1-{N-[(1R)-1-benzyl-2-((3S)-3-(dimethylaminomethyl)piperidin-1-yl)-2-oxoethyl]-N-methylcarbamoyl}-2-(2-naphthyl)ethyl)-N-methylamide




16

(2E)-5-Amino-5-methylhex-2-enoic acid N-((1R)-1-{N-[(1R)-1-benzyl-2-((2S)-2-((dimethylamino)methyl)pyrrolidin-1-yl)-2-oxoethyl]-N-methylcarbamoyl}-2-(2-naphthyl)ethyl)-N-methylamide



5


N-((1R)-1-{N-[(1R)-1-Benzyl-2-((2S)-2-((dimethylamino)methyl)pyrrolidin-1-yl)-2-oxoethyl]-N-methylcarbamoyl}-2-(2-naphthyl)ethyl)-N-methyl-3-((methylamino)methyl)benzamide




10

15


(2E)-5-Amino-5-methylhex-2-enoic acid N-((1R)-1-{N-[(1R)-1-benzyl-2-(4-(dimethylamino)piperidin-1-yl)-2-oxoethyl]-N-methylcarbamoyl}-2-(2-naphthyl)ethyl)-N-methylamide.




(2E)-5-Amino-5-methylhex-2-enoic acid N-methyl-N-[(1R)-1-(N-methyl-N-[(1R)-1-[N-methyl-N-(1-methylpiperidin-4-yl)carbamoyl]-2-phenylethyl]carbamoyl]-2-(2-naphthyl)ethyl]amide




- 5 3-Aminomethyl-N-((1R)-1-{N-[(1R)-1-benzyl-2-(4-methylpiperazin-1-yl)-2-oxoethyl]-N-methylcarbamoyl}-2-(2-naphthyl)ethyl)-N-methylbenzamide



- 10 (2E)-5-Amino-5-methylhex-2-enoic acid N-((1R)-1-{N-[(1R)-1-benzyl-2-(4-methylpiperazin-1-yl)-2-oxoethyl]-N-methylcarbamoyl}-2-(2-naphthyl)ethyl)-N-methylamide




(2E)-5-Amino-5-methylhex-2-enoic acid N-methyl-N-((1R)-1-{N-methyl-N-[(1R)-2-phenyl-1-((2,2,6,6-tetramethylpiperidin-4-yl)carbamoyl)ethyl]carbamoyl}-2-(2-naphthyl)ethyl)amide




5


3-Aminomethyl-N-methyl-N-((1R)-1-{N-methyl-N-[(1R)-2-phenyl-1-((2,2,6,6-tetramethylpiperidin-4-yl)carbamoyl)ethyl]carbamoyl}-2-(2-naphthyl)ethyl)benzamide




(2E)-5-Amino-3,5-dimethylhex-2-enoic acid N-methyl-N-((1R)-1-{N-methyl-N-[(1R)-2-phenyl-1-((2,2,6,6-tetramethylpiperidin-4-yl)carbamoyl)ethyl]carbamoyl}-2-(2-naphthyl)ethyl)amide




- 5 (2E)-4-(1-Aminocyclobutyl)but-2-enoic acid N-((1R)-1-{N-[(1R)-1-benzyl-2-(4-methylpiperazin-1-yl)-2-oxoethyl]-N-methylcarbamoyl}-2-(2-naphthyl)ethyl)-N-methylamide



(2E)-5-Amino-3,5-dimethylhex-2-enoic acid N-((1R)-1-{N-[(1R)-1-benzyl-2-(4-methylpiperazin-1-yl)-2-oxoethyl]-N-methylcarbamoyl}-2-(2-naphthyl)ethyl)-N-methylamide




(2E)-4-(1-Aminocyclobutyl)but-2-enoic acid N-((1R)-1-{N-[(1R)-1-benzyl-2-(4-hydroxypiperidin-1-yl)-2-oxoethyl]-N-methylcarbamoyl}-2-(biphenyl-4-yl)ethyl)-N-methylamide




5

(2E)-5-Amino-5-methylhex-2-enoic acid N-((1R)-1-{N-[(1R)-1-benzyl-2-(4-hydroxypiperidin-1-yl)-2-oxoethyl]-N-methylcarbamoyl}-2-(biphenyl-4-yl)ethyl)-N-methylamide




10

(2E)-5-Amino-3,5-dimethylhex-2-enoic acid N-((1R)-1-{N-[(1R)-1-benzyl-2-(4-hydroxypiperidin-1-yl)-2-oxoethyl]-N-methylcarbamoyl}-2-(biphenyl-4-yl)ethyl)-N-methylamide



5


(2E)-5-Amino-5-methylhex-2-enoic acid N-((1R)-1-{N-[(1R)-1-benzyl-2-(4-hydroxypiperidin-1-yl)-2-oxoethyl]-N-methylcarbamoyl}-2-(2-naphthyl)ethyl)-N-methylamide




10

15


(2E)-5-Amino-3,5-dimethylhex-2-enoic acid N-((1R)-1-{N-[(1R)-1-benzyl-2-(4-hydroxy-piperidin-1-yl)-2-oxoethyl]-N-methylcarbamoyl}-2-(2-naphthyl)ethyl)-N-methylamide




5 (2E)-4-(1-Aminocyclobutyl)but-2-enoic acid N-((1R)-1-{N-[(1R)-1-benzyl-2-(4-hydroxy-piperidin-1-yl)-2-oxoethyl]-N-methylcarbamoyl}-2-(2-naphthyl)ethyl)-N-methylamide




(2E)-5-Amino-5-methylhex-2-enoic acid N-((1R)-1-{N-[(1R)-1-(4-fluorobenzyl)-2-(4-hydroxy-piperidin-1-yl)-2-oxoethyl]-N-methylcarbamoyl}-2-(2-naphthyl)ethyl)-N-methylamide




(2E)-5-Amino-3,5-dimethylhex-2-enoic acid N-((1R)-1-{N-[(1R)-1-(4-fluorobenzyl)-2-(4-hydroxypiperidin-1-yl)-2-oxoethyl]-N-methylcarbamoyl}-2-(2-naphthyl)ethyl)-N-methylamide




5 (2E)-5-Amino-5-methylhex-2-enoic acid N-((1R)-1-{N-[(1R)-1-benzyl-2-(4-hydroxy-4-(2-thienyl)piperidin-1-yl)-2-oxoethyl]-N-methylcarbamoyl}-2-(2-naphthyl)ethyl)-N-methylamide




(2E)-5-Amino-5-methylhex-2-enoic acid N-((1R)-1-{N-[(1R)-1-(3-hydroxycyclohexylcarbamoyl)-2-phenylethyl]-N-methylcarbamoyl}-2-(2-naphthyl)ethyl)-N-methylamide




(2E)4-(1-Aminocyclobutyl)but-2-enoic acid N-((1R)-1-{N-[(1R)-1-benzyl-2-(4-(dimethylamino)piperidin-1-yl)-2-oxoethyl]-N-methylcarbamoyl}-2-(2-naphthyl)ethyl)-N-methylamide




(2E)-5-Amino-5-methylhex-2-enoic acid N-((1R)-1-{N-[(2R)-2-(4-hydroxypiperidin-1-yl)-2-oxo-1-((2-thienyl)methyl)ethyl]-N-methylcarbamoyl}-2-(2-naphthyl)ethyl)-N-methylamide



(2E)-5-Amino-3,5-dimethylhex-2-enoic acid N-((1R)-1-{N-[(2R)-2-(4-hydroxypiperidin-1-yl)-2-oxo-1-((2-thienyl)methyl)ethyl]-N-methylcarbamoyl}-2-(2-naphthyl)ethyl)-N-methylamide




(2E)-5-Amino-5-methylhex-2-enoic acid N-((1R)-2-(biphenyl-4-yl)-1-{N-[(2R)-2-(4-hydroxypiperidin-1-yl)-2-oxo-1-((2-thienyl)methyl)ethyl]-N-methylcarbamoyl}ethyl)-N-methylamide



5


(2E)-5-Amino-3,5-dimethylhex-2-enoic acid N-((1R)-2-(biphenyl-4-yl)-1-{N-[(1R)-2-(4-hydroxypiperidin-1-yl)-2-oxo-1-((2-thienyl)methyl)ethyl]-N-methylcarbamoyl}ethyl)-N-methylamide

10




26

(2E)-5-Methyl-5-(methylamino)hex-2-enoic acid N-((1R)-1-{N-[(1R)-1-benzyl-2-(4-hydroxypiperidin-1-yl)-2-oxoethyl]-N-methylcarbamoyl}-2-(biphenyl-4-yl)ethyl)-N-methylamide



5

(2E)-4-(1-Aminocyclobutyl)but-2-enoic acid ((1R)-1-{N-[(1R)-1-benzyl-2-(4-hydroxypiperidin-1-yl)-2-oxoethyl]-N-methylcarbamoyl}-2-(biphenyl-4-yl)ethyl)amide




and pharmaceutically acceptable salts thereof.

10


General Methods

The methods illustrated in below schemes I - III are by no mean intended to limit the present invention in any aspect, but should only be seen as a guidance for how the present compounds may be prepared.

Scheme I




Amines of type




- 5 may be synthesized from a BOC-protected amino acid (cf. scheme I). The acid is transformed into an ester by reaction with or without a reagent such as e. g. 1-hydroxybenzotriazole, 1-hydroxy-7-azabenzotriazole, or 3-hydroxy-1,2,3-benzotriazole-4(3H)-one and a reagent such e.g. as N-(3-dimethylaminopropyl)-N'-ethylcarbodiimide hydrochloride or diisopropylcarbodiimide and a catalyst such as N,N-dimethylaminopyridine. The ester may be reduced by a suitable reagent, such as diisobutylaluminum hydride in a appropriate solvent such as e.g. toluene, dichloromethane, ether, or tetrahydrofuran to give the BOC-protected aldehyde or alcohol. If the alcohol is obtained, the alcohol may be oxidized to the corresponding aldehyde, by a suitable method, such as e.g. dimethylsulfoxide/oxalyl chloride/ triethylamine or dimethylsulfoxide/sulfotrioxide/pyridine, pyridinium dichromate, or pyridinium chlorochromate. A reductive amination with an appropriate amine $N(R^{15})(R^{16})$ and a suitable reagent, such as sodium cyanoborohydride or sodium triacetoxyborohydride in a suitable solvent such as e.g. alcohols may yield the BOC-protected amine. If at least one of R^{15} or R^{16} is hydrogen, the amino group may be protected by a method known to those skilled in the art and described in the literature as e.g. T. W. Greene, P. G. M. Wuts Protective groups in organic synthesis, 2nd edition, Wiley, New York, before carrying out the following steps. The removal of the BOC-protection group can be achieved by a method known to those skilled in the art as described in T. W. Greene, P. G. M. Wuts Protective groups in organic synthesis, 2nd edition, Wiley, New York, such as e.g. hydrogen chloride in ethyl acetate, or trifluoroacetic acid in dichloromethane.

Scheme II




Compounds of the type of formula I may be synthesized by coupling of an amine of type




and a suitable protected acid with or without a coupling reagent such as e.g. 1-hydroxy-
 5 benzotriazole, 1-hydroxy-7-azabenzotriazole, or 3-hydroxy-1,2,3-benzotriazole-4(3H)-one
 and a reagent such e.g. as N-(3-dimethylaminopropyl)-N'-ethylcarbodiimide hydrochloride or
 diisopropylcarbodiimide in a suitable solvent, such as N,N-dimethylformamide or dichloro-
 methane (cf. scheme II). The product may be deprotected at the nitrogen of the acid by a
 method known for a person skilled in the art and described in the literature e.g. in T. W.

10 Greene, P. G. M. Wuts Protective groups in organic synthesis, 2nd edition, Wiley, New York.
 The product is coupled with a suitable protected acid with or without a coupling reagent such
 as e.g. 1-hydroxybenzotriazole, 1-hydroxy-7-azabenzotriazole, or 3-hydroxy-1,2,3-benzo-
 triazole-4(3H)-one and a reagent such e.g. as N-(3-dimethylaminopropyl)-N'-ethyl-
 carbodiimide hydrochloride or diisopropylcarbodiimide in a suitable solvent, such as N,N-
 15 dimethylformamide or dichloromethane. The product may be deprotected at the nitrogen of
 the acid by a method known for a person skilled in the art and described in the literature e.g.
 in T. W. Greene, P. G. M. Wuts Protective groups in organic synthesis, 2nd edition, Wiley,
 New York. The product is coupled with suitable protected acid with or without a coupling
 reagent such as e.g. 1-hydroxybenzotriazole, 1-hydroxy-7-azabenzotriazole, or 3-hydroxy-
 20 1,2,3-benzotriazole-4(3H)-one and a reagent such e.g. as N-(3-dimethylaminopropyl)-N'-
 ethylcarbodiimide hydrochloride or diisopropylcarbodiimide in a suitable solvent, such as
 N,N-dimethylformamide or dichloromethane. All protection groups may be removed by a
 method known for a person skilled in the art and described in the literature e.g. in T. W.
 Greene, P. G. M. Wuts Protective groups in organic synthesis, 2nd edition, Wiley, New York.

Scheme III



Compounds of the type of formula I may be synthesized by coupling of an amine of type



5

and a suitable protected acid with or without a coupling reagent such as e.g. 1-hydroxybenzotriazole, 1-hydroxy-7-azabenzotriazole, or 3-hydroxy-1,2,3-benzotriazole-4(3H)-one and a reagent such e.g. as N-(3-dimethylaminopropyl)-N'-ethylcarbodiimide hydrochloride or diisopropylcarbodiimide in a suitable solvent, such as N,N-dimethylformamide or dichloromethane (cf. scheme III). The product may be deprotected at the nitrogen of the acid by a method known for a person skilled in the art and described in the literature e.g. in T. W. Greene, P. G. M. Wuts Protective groups in organic synthesis, 2nd edition, Wiley, New York. The product is coupled with a suitable protected acid with or without a coupling reagent such as e.g. 1-hydroxybenzotriazole, 1-hydroxy-7-azabenzotriazole, or 3-hydroxy-1,2,3-benzotriazole-4(3H)-one and a reagent such e.g. as N-(3-dimethylaminopropyl)-N'-ethylcarbodiimide hydrochloride or diisopropylcarbodiimide in a suitable solvent, such as N,N-dimethylformamide or dichloromethane. The product may be deprotected at the nitrogen of the acid by a method known for a person skilled in the art and described in the literature e.g. in T. W. Greene, P. G. M. Wuts Protective groups in organic synthesis, 2nd edition, Wiley, New York. The product is coupled with suitable protected acid with or without a coupling reagent such as e.g. 1-hydroxybenzotriazole, 1-hydroxy-7-azabenzotriazole, or 3-hydroxy-1,2,3-benzotriazole-4(3H)-one and a reagent such e.g. as N-(3-dimethylaminopropyl)-N'-ethylcarbodiimide hydrochloride or diisopropylcarbodiimide in a suitable solvent, such as N,N-dimethylformamide or dichloromethane. All protection groups may be removed by a method known for a person skilled in the art and described in the literature e.g. in T. W. Greene, P. G. M. Wuts Protective groups in organic synthesis, 2nd edition, Wiley, New York.

The compounds of formula I exhibit an improved resistance to proteolytic degradation by enzymes because they are non-natural, in particular because the natural amide bonds are replaced by non-natural amide bond mimetics. The increased resistance to proteolytic degradation of the compounds of the invention in comparison with known hormone releasing peptides is expected to improve their bioavailability compared to that of the peptides suggested in the prior literature.

In the above structural formulas and throughout the present specification, the following terms have the indicated meanings:

The C₁₋₆-alkyl, C₁₋₆-alkylene, C₁₋₄-alkyl or C₁₋₄-alkylene groups specified above are intended to include those alkyl or alkylene groups of the designated length in either a linear or branched or cyclic configuration. Examples of linear alkyl are methyl, ethyl, propyl, butyl, pentyl, and hexyl and their corresponding divalent moieties, such as ethylene. Examples of branched alkyl are isopropyl, sec-butyl, tert-butyl, isopentyl, and isohexyl and their corresponding divalent moieties, such as isopropylene. Examples of cyclic alkyl are C₃₋₆-cycloalkyl such as cyclopropyl, cyclobutyl, cyclopentyl and cyclohexyl and their corresponding divalent moieties, such as cyclopropylene.

The C₁₋₆-alkoxy groups specified above are intended to include those alkoxy groups of the designated length in either a linear or branched or cyclic configuration. Examples of linear alkoxy are methoxy, ethoxy, propoxy, butoxy, pentoxy, and hexoxy. Examples of branched alkoxy are isopropoxy, sec-butoxy, tert-butoxy, isopentoxy, and isohexoxy. Examples of cyclic alkoxy are C₃₋₆-cycloalkoxy such as cyclopropyloxy, cyclobutyloxy, cyclopentyloxy and cyclohexyloxy.

In the present context, the term "aryl" is intended to include monovalent carbocyclic aromatic ring moieties, being either monocyclic, bicyclic or polycyclic, e.g. selected from the group consisting of phenyl and naphthyl, optionally substituted with one or more C₁₋₆-alkyl, C₁₋₆-alkoxy, halogen, amino or aryl.

In the present context, the term "arylene" is intended to include divalent carbocyclic aromatic ring moieties, being either monocyclic, bicyclic or polycyclic, e.g. selected from the group con-

sisting of phenylene and naphthylene, optionally substituted with one or more C₁₋₆-alkyl, C₁₋₆-alkoxy, halogen, amino or aryl.

In the present context, the term "hetaryl" is intended to include monovalent heterocyclic aro-

5 matic ring moieties, being either monocyclic, bicyclic or polycyclic, e.g. selected from the group consisting of pyridyl, 1-H-tetrazol-5-yl, thiazolyl, imidazolyl, indolyl, pyrimidinyl, thiadiazolyl, pyrazolyl, oxazolyl, isoxazolyl, oxadiazolyl, thienyl, quinoliny, pyrazinyl, or isothiazolyl, optionally substituted with one or more C₁₋₆-alkyl, C₁₋₆-alkoxy, halogen, amino or aryl.

10 In the present context, the term "hetarylene" is intended to include divalent heterocyclic aromatic ring moieties, being either monocyclic, bicyclic or polycyclic, e.g. selected from the group consisting of pyridinediyl, 1-H-tetrazolediyl, thiazoldiyl, imidazolediyl, indolediyl, pyrimidinediyl, thiadiazolelediyl, pyrazolelediyl, oxazolelediyl, isoxazolelediyl, oxadiazolelediyl, thiophenediyl, quinolinediyl, pyrazinediyl, or isothiazolelediyl, optionally substituted with one or more C₁₋₆-alkyl, C₁₋₆-
15 alkoxy, halogen, amino or aryl.

In the present context, the term "heterocyclic system" is intended to include aromatic as well as non-aromatic ring moieties, which may be monocyclic, bicyclic or polycyclic, and contain in their ring structure at least one, such as one, two or three, nitrogen atom(s), and optionally one
20 or more, such as one or two, other hetero atoms, e.g. sulphur or oxygen atoms. The heterocyclic system is preferably selected from pyrazole, pyridazine, triazine, indazole, phthalazine, cinnoline, pyrazolidine, pyrazoline, aziridine, dithiazine, pyrrol, imidazol, pyrazole, isoindole, indole, indazole, purine, pyrrolidine, pyrroline, imidazolidine, imidazoline, pyrazolidine, pyrazoline, piperidine, piperazine, indoline, isoindoline, or morpholine, optionally substituted
25 with one or more C₁₋₆-alkyl, C₁₋₆-alkoxy, halogen, amino, oxy or aryl.

The term "halogen" is intended to include chlorine (Cl), fluorine (F), bromine (Br) and iodine (I).

In the context of the present application, the term "growth hormone secretagogue" is intended
30 to include any compound which has the capacity, directly or indirectly, of inducing (i.e. stimulating or increasing) the release of growth hormone from the pituitary gland. The term "growth hormone secretagogue" includes growth hormone releasing peptides, growth hormone releasing peptidomimetics, and growth hormone releasing compounds of a nonpeptidyl nature.

The compounds of the present invention may optionally be on a pharmaceutically acceptable salt form such as the pharmaceutically acceptable acid addition salts of compounds of formula I which include those prepared by reacting the compound of formula I with an inorganic or organic acid such as hydrochloric, hydrobromic, sulfuric, acetic, phosphoric, lactic, malic, maleic, mandelic phthalic, citric, glutaric, gluconic, methanesulfonic, salicylic, succinic, tartaric, toluenesulfonic, trifluoroacetic, sulfamic or fumaric acid and/or water.

The compounds of formula I may be administered in pharmaceutically acceptable acid addition salt form or, where appropriate, as a alkali metal or alkaline earth metal or lower alkylammonium salt. Such salt forms are believed to exhibit approximately the same order of activity as the free base forms.

In another aspect, the present invention relates to a pharmaceutical composition comprising, as an active ingredient, a compound of the general formula I or a pharmaceutically acceptable salt thereof together with a pharmaceutically acceptable carrier or diluent.

Pharmaceutical compositions containing a compound of the present invention may be prepared by conventional techniques, e.g. as described in Remington's Pharmaceutical Sciences, 1985 or in Remington: The Science and Practice of Pharmacy, 19th Edition (1995). The compositions may appear in conventional forms, for example capsules, tablets, aerosols, solutions, suspensions or topical applications.

The pharmaceutical carrier or diluent employed may be a conventional solid or liquid carrier. Examples of solid carriers are lactose, terra alba, sucrose, cyclodextrin, talc, gelatin, agar, pectin, acacia, magnesium stearate, stearic acid or lower alkyl ethers of cellulose. Examples of liquid carriers are syrup, peanut oil, olive oil, phospholipids, fatty acids, fatty acid amines, polyoxyethylene or water.

Similarly, the carrier or diluent may include any sustained release material known in the art, such as glyceryl monostearate or glyceryl distearate, alone or mixed with a wax.

If a solid carrier is used for oral administration, the preparation may be tableted, placed in a hard gelatin capsule in powder or pellet form or it can be in the form of a troche or lozenge. The amount of solid carrier will vary widely but will usually be from about 25 mg to about 1 g. If a liquid carrier is used, the preparation may be in the form of a syrup, emulsion, soft gelatin capsule or sterile injectable liquid such as an aqueous or non-aqueous liquid suspension or solution.

A typical tablet which may be prepared by conventional tableting techniques may contain:

10 Core:

Active compound (as free compound or salt thereof)	10mg
Colloidal silicon dioxide (Aerosil)	1.5mg
Cellulose, microcryst. (Avicel)	70mg
Modified cellulose gum (Ac-Di-Sol)	7.5mg

15 Magnesium stearate

Coating:

HPMC approx.	9mg
*Mywacett 9-40 T approx.	0.9mg

20

*Acylated monoglyceride used as plasticizer for film coating.

For nasal administration, the preparation may contain a compound of formula I dissolved or suspended in a liquid carrier, in particular an aqueous carrier, for aerosol application. The carrier may contain additives such as solubilizing agents, e.g. propylene glycol, surfactants, absorption enhancers such as lecithin (phosphatidylcholine) or cyclodextrin, or preservatives such as parabenes.

It has been demonstrated that compounds of the general formula I possess the ability to release endogenous growth hormone *in vivo*. The compounds may therefore be used in the treatment of conditions which require increased plasma growth hormone levels such as in growth hormone deficient humans or in elderly patients or livestock.

Thus, in a particular aspect, the present invention relates to a pharmaceutical composition for stimulating the release of growth hormone from the pituitary, the composition comprising, as an active ingredient, a compound of the general formula I or a pharmaceutically acceptable salt thereof together with a pharmaceutically acceptable carrier or diluent.

5

In a further aspect, the present invention relates to a method of stimulating the release of growth hormone from the pituitary, the method comprising administering to a subject in need thereof an effective amount of a compound of the general formula I or a pharmaceutically acceptable salt thereof.

10

In a still further aspect, the present invention relates to a method of treating growth retardation in connection with asthma, the method comprising administering to a subject in need thereof an effective amount of a growth hormone secretagogue or a pharmaceutically acceptable salt thereof. In a particular embodiment, the present invention relates to a method of treating growth retardation in connection with asthma, the method comprising administering to a subject in need thereof an effective amount of a compound of the general formula I or a pharmaceutically acceptable salt thereof.

15

In a still further aspect, the present invention relates to a method of treating growth retardation in connection with juvenile rheumatic arthritis or cystic fibrosis, the method comprising administering to a subject in need thereof an effective amount of a growth hormone secretagogue or a pharmaceutically acceptable salt thereof. In one embodiment, the present invention relates to a method of treating growth retardation in connection with juvenile rheumatic arthritis, the method comprising administering to a subject in need thereof an effective amount of a growth hormone secretagogue or a pharmaceutically acceptable salt thereof. In a second embodiment, the present invention relates to a method of treating growth retardation in connection with cystic fibrosis, the method comprising administering to a subject in need thereof an effective amount of a growth hormone secretagogue or a pharmaceutically acceptable salt thereof. In a particular embodiment, the present invention relates to a method of treating growth retardation in connection with juvenile rheumatic arthritis, the method comprising administering to a subject in need thereof an effective amount of a compound of the general formula I or a pharmaceutically acceptable salt thereof. In another particular embodiment, the present invention relates to a method of treating growth retardation in connection with cystic fibrosis, the method

20

25

30

comprising administering to a subject in need thereof an effective amount of a compound of the general formula I or a pharmaceutically acceptable salt thereof.

In a still further aspect, the present invention relates to the use of a compound of the general formula I or a pharmaceutically acceptable salt thereof for the preparation of a medicament for stimulating the release of growth hormone from the pituitary.

To those skilled in the art, it is well known that the current and potential uses of growth hormone in humans are varied and multitudinous. Thus, compounds of formula I can be administered for purposes stimulating release of growth hormone from the pituitary and would then have similar effects or uses as growth hormone itself. Compounds of formula I are useful for: stimulation of growth hormone release in the elderly, prevention of catabolic side effects of glucocorticoids, prevention and treatment of osteoporosis, treatment of chronic fatigue syndrome (CFS), treatment of acute fatigue syndrome and muscle loss following elective surgery, stimulation of the immune system, acceleration of wound healing, accelerating bone fracture repair, accelerating complicated fractures, e.g. distraction osteogenesis, treatment of wasting secondary to fractures, treatment of growth retardation, treating growth retardation resulting from renal failure or insufficiency, treatment of cardiomyopathy, treatment of wasting in connection with chronic liver disease, treatment of thrombocytopenia, treatment of growth retardation in connection with Crohn's disease, treatment of short bowel syndrome, treatment of wasting in connection with chronic obstructive pulmonary disease (COPD), treatment of complications associated with transplantation, treatment of physiological short stature including growth hormone deficient children and short stature associated with chronic illness, treatment of obesity and growth retardation associated with obesity, treatment of anorexia, treatment of growth retardation associated with the Prader-Willi syndrome and Turner's syndrome; increasing the growth rate of a patient having partial growth hormone insensitive syndrome, accelerating the recovery and reducing hospitalization of burn patients; treatment of intrauterine growth retardation, skeletal dysplasia, hypercortisolism and Cushing's syndrome; induction of pulsatile growth hormone release; replacement of growth hormone in stressed patients, treatment of osteochondrodysplasias, Noonan's syndrome, schizophrenia, depressions, Alzheimer's disease, delayed wound healing and psychosocial deprivation, treatment of catabolism in connection with pulmonary dysfunction and ventilator dependency; treatment of cardiac failure or related vascular dysfunction, treatment of impaired cardiac function, treatment or prevention of myo-

cardial infarction, lowering blood pressure, protection against ventricular dysfunction or prevention of reperfusion events; treatment of adults in chronic dialysis; attenuation of protein catabolic responses after major surgery, reducing cachexia and protein loss due to chronic illness such as cancer or AIDS; treatment of hyperinsulinemia including nesidioblastosis, adjuvant treatment for ovulation induction; stimulation of thymic development and prevention of the age-related decline of thymic function, treatment of immunosuppressed patients; treatments of sarcopenia, treatment of wasting in connection with AIDS; improvement in muscle strength, mobility, maintenance of skin thickness, treatment of metabolic homeostasis and renal homeostasis in the frail elderly, stimulation of osteoblasts, bone remodelling and cartilage growth; regulation of food intake; stimulation of the immune system in companion animals and treatment of disorder of aging in companion animals, promoting growth in livestock and stimulation of wool growth in sheep, increasing milk production in livestock, treatment of metabolic syndrome (syndrome X), treatment of insulin resistance, including NIDDM, in mammals, e.g. humans, treatment of insulin resistance in the heart, improvement of sleep quality and correction of the relative hyposomatotropism of senescence due to high increase in REM sleep and a decrease in REM latency, treatment of hypothermia, treatment of frailty associated with aging, treatment of congestive heart failure, treatment of hip fractures, treatment of immune deficiency in individuals with a depressed T4/T8 cell ratio, treatment of muscular atrophy, treatment of musculoskeletal impairment in elderly, enhancing the activity of protein kinase B (PKB), improvement of the overall pulmonary function, and treatment of sleep disorders.

Within the context of the present application, the term "treatment" is also intended to include prophylactic treatment.

In a further aspect the present invention relates to the use of a growth hormone secretagogue or a pharmaceutically acceptable salt thereof for the preparation of a medicament for the treatment of growth retardation in connection with asthma. In one particular embodiment the invention relates to the use of a compound of the general formula I or a pharmaceutically acceptable salt thereof for the treatment of growth retardation in connection with asthma. In a second particular embodiment the invention relates to the use of growth hormone releasing peptides, growth hormone releasing peptidomimetics, or growth hormone releasing compounds of a nonpeptidyl nature or a pharmaceutically acceptable salt thereof for the treatment of growth retardation in connection with asthma.

In a still further aspect the present invention relates to the use of a growth hormone secretagogue or a pharmaceutically acceptable salt thereof for the preparation of a medicament for the treatment of growth retardation in connection with juvenile rheumatic arthritis or systic fibrosis. In one embodiment the invention relates to the use of a growth hormone secretagogue or a pharmaceutically acceptable salt thereof for the preparation of a medicament for the treatment of growth retardation in connection with juvenile rheumatic arthritis. In a second embodiment the invention relates to the use of a growth hormone secretagogue or a pharmaceutically acceptable salt thereof for the preparation of a medicament for the treatment of growth retardation in connection with systic fibrosis. In one particular embodiment the invention relates to the use of a compound of the general formula I or a pharmaceutically acceptable salt thereof for the treatment of growth retardation in connection with juvenile rheumatic arthritis. In another particular embodiment the invention relates to the use of a compound of the general formula I or a pharmaceutically acceptable salt thereof for the treatment of growth retardation in connection with systic fibrosis. In a further particular embodiment the invention relates to the use of growth hormone releasing peptides, growth hormone releasing peptidomimetics, or growth hormone releasing compounds of a nonpeptidyl nature or a pharmaceutically acceptable salt thereof for the treatment of growth retardation in connection with juvenile rheumatic arthritis. In a still further particular embodiment the invention relates to the use of growth hormone releasing peptides, growth hormone releasing peptidomimetics, or growth hormone releasing compounds of a nonpeptidyl nature or a pharmaceutically acceptable salt thereof for the treatment of growth retardation in connection with systic fibrosis.

For the above indications the dosage will vary depending on the growth hormone secretagogue employed, e.g. on the compound of formula I employed, on the mode of administration and on the therapy desired. However, generally dosage levels between 0.0001 and 100 mg/kg body weight daily are administered to patients and animals to obtain effective release of endogenous growth hormone. Moreover the compounds of formula I have no or substantially no side-effects, when administered in the above dosage levels, such side-effects being e.g. release of LH, FSH, TSH, ACTH, vasopressin, oxytocin, cortisol and/or prolactin. Usually, dosage forms suitable for oral, nasal, pulmonal or transdermal administration comprise from about 0.0001 mg to

about 100 mg, preferably from about 0.001 mg to about 50 mg of the compounds of formula I admixed with a pharmaceutically acceptable carrier or diluent.

The dosage of the compounds according to this invention is suitably 0.01-500 mg/day, e.g.

5 from about 5 to about 50 mg, such as about 10 mg per dose, when administered to patients, e.g. humans, as a drug.

Optionally, the pharmaceutical composition of the invention may comprise a compound of formula I combined with one or more compounds exhibiting a different activity, e.g., an antibiotic
10 or other pharmacologically active material.

The route of administration may be any route which effectively transports the active compound to the appropriate or desired site of action, such as oral, nasal, pulmonary, transdermal or par-
enteral, the oral route being preferred.

15

Apart from the pharmaceutical use of the compounds of formula I, they may be useful in vitro tools for investigating the regulation of growth hormone release.

Compounds of formula I may also be useful in vivo tools for evaluating the growth hormone
20 releasing capability of the pituitary. For example, serum samples taken before and after administration of these compounds to humans can be assayed for growth hormone. Comparison of the growth hormone in each serum sample would directly determine the ability of the patients pituitary to release growth hormone.

25 Compounds of formula I may be administered to commercially important animals to increase their rate and extent of growth, and to increase milk and wool production.

A further use of growth hormone secretagogue compounds of formula I is in combination with other secretagogues such as GHRP (2 or 6), GHRH and its analogues, growth hormone and
30 its analogues or somatomedins including IGF-1 and IGF-2.

Pharmacological Methods

Compounds of formula I may be evaluated in vitro for their efficacy and potency to release growth hormone in rat pituitary primary cultures, and such evaluation may be performed as described below.

The isolation of rat pituitary cells is a modification of O. Sartor et al., *Endocrinology* **116**, 1985, pp. 952-957. Male albino Sprague-Dawley rats (250 +/- 25 grams) were purchased from Møllegaard, Lille Skensved, Denmark. The rats were housed in group cages (four animals/cage) and placed in rooms with 12 hour light cycle. The room temperature varied from 19-24°C and the humidity from 30 - 60%.

The rats were decapitated and the pituitaries dissected. The neurointermediate lobes were removed and the remaining tissue was immediately placed in icecold isolation buffer (Gey's medium (Gibco 041-04030) supplemented with 0.25% D-glucose, 2% non-essential amino acids (Gibco 043-01140) and 1% bovine serum albumine (BSA) (Sigma A-4503)). The tissue was cut into small pieces and transferred to isolation buffer supplemented with 3.8 mg/ml of trypsin (Worthington #3707 TRL-3) and 330 mg/ml of DNase (Sigma D-4527). This mixture was incubated at 70 rotations/min for 35 min at 37°C in a 95/5% atmosphere of O₂/CO₂. The tissue was then washed three times in the above buffer. Using a standard pasteur pipette, the tissue was then aspirated into single cells. After dispersion, cells were filtered through a nylon filter (160 µm) to remove undigested tissue. The cell suspension was washed 3 times with isolation buffer supplemented with trypsin inhibitor (0.75 mg/ml, Worthington #2829) and finally resuspended in culture medium; DMEM (Gibco 041-01965) supplemented with 25 mM HEPES (Sigma H-3375), 4 mM glutamine (Gibco 043-05030H), 0.075% sodium bicarbonate (Sigma S-8875), 0.1% non-essential amino acid, 2.5% fetal calf serum (FCS, Gibco 011-06290), 3% horse serum (Gibco 034-06050), 10% fresh rat serum, 1 nM T₃ (Sigma T-2752) and 40 mg/l dexamethasone (Sigma D-4902) pH 7.3, to a density of 2 x 10⁵ cells/ml. The cells were seeded into microtiter plates (Nunc, Denmark), 200 µl/well, and cultured for 3 days at 37°C and 8% CO₂.

Compound testing

After culturing, the cells were washed twice with stimulation buffer (Hanks Balanced Salt Solution (Gibco 041-04020) supplemented with 1% BSA (Sigma A-4503), 0.25% D-glucose (Sigma G-5250) and 25 mM HEPES (Sigma H-3375) pH 7.3) and preincubated for 1 hour at 37°C. The buffer was exchanged with 90 ml stimulation buffer (37°C). Ten ml test compound solution was added and the plates were incubated for 15 min at 37°C and 5% CO₂. The medium was decanted and analyzed for GH content in an rGH SPA test system.

All compounds were tested in doses ranging from 10 pM to 100 mM. A dose-response relation was constructed using the Hill equation (Fig P, Biosoft). The efficacy (maximal GH released, E_{max}) was expressed in % of the E_{max} of GHRP-6. The potency (EC₅₀) was determined as the concentration inducing half maximal stimulation of the GH release.

Compounds of formula I may be evaluated for their metabolic stability using the procedure described below:

Compounds is dissolved at a concentration of 1 mg/ml in water. 25 ml of this solution is added to 175 ml of the respective enzyme-solution (resulting in an enzyme:substrate ratio (w/w) of approximately 1:5). The solution is left at 37°C overnight. 10 ml of the various degradation solutions is analyzed against a corresponding zero-sample using flow injection electrospray mass spectrometry (ESMS) with selected ion monitoring of the molecular ion. If the signal has decreased more than 20% compared to the zero-sample, the remainder of the solution is analyzed by HPLC and mass spectrometry in order to identify the extent and site(s) of degradation precisely.

Several standard peptides (ACTH 4-10, Angiotensin 1-14 and Glucagon) have been included in the stability tests in order to verify the ability of the various solutions to degrade peptides.

Standard peptides (angiotensin 1-14, ACTH 4-10 and glucagon) were purchased from Sigma, MO, USA)

Enzymes (trypsin, chymotrypsin, elastase aminopeptidase M and carboxypeptidase Y and B) were all purchased from Boehringer Mannheim GmbH (Mannheim, Germany)

5 Pancreatic enzyme mix: trypsin, chymotrypsin and elastase in 100 mM ammoniumbicarbonate pH 8.0 (all concentrations 0.025 mg/ml).

Carboxypeptidase mix: carboxypeptidase Y and B in 50 mM ammoniumacetate pH 4.5 (all concentrations 0.025 mg/ml).

10 Aminopeptidase M solution: aminopeptidase M (0.025 mg/ml) in 100 mM ammoniumbicarbonate pH 8.0

Mass spectrometric analysis was performed using two different mass spectrometers. A Sciex API III triple quadrupole LC-MS instrument (Sciex instruments, Thornhill, Ontario) equipped
15 with an electrospray ion-source and a Bio-Ion 20 time-of-flight Plasma Desorption instrument (Bio-Ion Nordic AB, Uppsala, Sweden).

Quantification of the compounds (before and after degradation) was done on the API III instrument using single ion monitoring of the molecular ion in question with flow injection of the
20 analyte. The liquid flow (MeOH:water 1:1) of 100 ml/min was controlled by an ABI 140B HPLC unit (Perkin-Elmer Applied Biosystems Divisions, Foster City, CA). The instrument parameters were set to standard operation conditions, and SIM monitoring was performed using the most intense molecular ion (in most cases this corresponded to the doubly charged molecular ion).

25 Identification of degradation products furthermore involved the use of plasma desorption mass spectrometry (PDMS) with sample application on nitrocellulose coated targets and standard instrumental settings. The accuracy of the hereby determined masses is generally better than 0.1%.

30 Separation and isolation of degradation products was done using a HY-TACH C-18 reverse phase 4.6x105 mm HPLC column (Hewlett-Packard Company, Palo Alto, CA) with a standard acetonitril: TFA separation gradient. The HPLC system used was HP1090M (Hewlett-Packard Company, Palo Alto, CA).

Peptide derivative	MW/SIM ion (amu)	Carboxy-peptidase mix	Pan. enzyme mix
Standards			
ACTH 4-10	1124.5/562.8	+	-
Glucagon	3483/871.8	-	-
Insulin (B23-29)	859.1/430.6		
Angiotensin 1-14	1760.1/881.0	-	-
GHRP-2	817.4/409.6	-	-
GHRP-6	872.6/437.4	-	-

+: Stable (less than 20% decrease in SIM signal after 24 h in degradation solution)

-: Unstable (more than 20% decrease in SIM signal after 24 h in degradation solution)

5

Any novel feature or combination of features described herein is considered essential to this invention.

EXAMPLES:

The process for preparing compounds of formula I and preparations containing them is further illustrated in the following examples, which however, are not to be construed as limiting.

5

The structures of the compounds are confirmed by either elemental analysis (MA) nuclear magnetic resonance (NMR) or mass spectrometry (MS). NMR shifts (δ) are given in parts per million (ppm) and only selected peaks are given. mp is melting point and is given in °C. Column chromatography was carried out using the technique described by W.C. Still et al, J. Org. Chem. 1978, 43, 2923-2925 on silica gel 60. Compounds used as starting materials are either known compounds or compounds which can readily be prepared by methods known per se.

10

HPLC-Analysis:

15

Method A1.

The RP-analysis was performed using UV detections at 214, 254, 276, and 301 nm on a 218TP54 4.6 mm x 250 mm 5m C-18 silica column (The Separations Group, Hesperia), which was eluted at 1 mL/min at 42°C. The column was equilibrated with 5% acetonitrile in a buffer consisting of 0.1 M ammonium sulfate, which was adjusted to pH 2.5 with 4M sulfuric acid. after injection the sample was eluted by a gradient of 5% to 60% acetonitrile in the same buffer during 50 min.

20

Method B1.

The RP-analysis was performed using UV detections at 214, 254, 276, and 301 nm on a 218TP54 4.6 mm x 250 mm 5m C-18 silica column (The Separations Group, Hesperia), which was eluted at 1 mL/min at 42°C. The column was equilibrated with 5% (acetonitrile + 0.1 % TFA) in an aqueous solution of TFA in water (0.1%). After injection the sample was eluted by a gradient of 5% to 60% (acetonitrile + 0.1% TFA) in the same aqueous buffer during 50 min.

25

30

Abbreviations:

TLC: thin layer chromatography

DMSO: dimethylsulfoxide

min: minutes

5 h: hours

Boc: tert butyloxycarbonyl

DMF: dimethylformamide

THF: tetrahydrofuran

EDAC: N-ethyl-N'-dimethylaminopropylcarbodiimide hydrochloride

10 HOAt: 1-hydroxy-7-azabenzotriazole

DIEA: diisopropylethylamine

TFA: trifluoroacetic acid


Buildingblocks:

15

N-methylated aminoacids used in the following examples were prepared as in Can. J. Chem. 1977, 55, 906.

3-Hydroxy-1,1-dimethylpropylcarbamic acid tert-butyl ester:

20




At 0 °C, ethyl chloroformate (1.10 mL, 11.5 mmol) was given dropwise to a solution of 3-tert-butoxycarbonylamino-3-methylbutanoic acid (2.50 g, 11.5 mmol) and triethylamine (1.92 mL, 13.8 mmol) in tetrahydrofuran (10 mL). The solution was stirred for 40 min at 0 °C. The formed precipitate was filtered off and washed with tetrahydrofuran (20 mL). The liquid was immediately cooled to 0 °C. A 2M solution of lithium boronhydride in tetrahydrofuran (14.4 mL, 28.8 mmol) was added dropwise. The solution was stirred at 0 °C for 2 h, and then warmed to room temperature. over a period of 4 h. It was cooled to 0 °C. Methanol (5 mL) was added carefully. 1N Hydrochloric acid (100 mL) was added. The solution was extracted with ethyl acetate (2 x 100 mL, 3 x 50 mL). The combined organic layers were washed with saturated sodium

hydrogen carbonate solution (100 mL) and dried over magnesium sulfate. The solvent was removed in vacuo. The crude product was chromatographed on silica (110 g) with ethyl acetate/heptane 1:2 to give 1.84 g of 3-hydroxy-1,1-dimethylpropylcarbamic acid tert-butyl ester.

- 5 $^1\text{H-NMR}$ (CDCl_3): d 1.33 (s, 6 H); 1.44 (s, 9 H); 1.88 (t, 2 H); 1.94 (br, 1 H); 3.75 (q, 2 H); 4.98 (br, 1 H).


3-(tert-Butoxycarbonylamino)-3-methylbutanal:

10



- DMSO (1.22 mL, 17.2 mmol) was added to a solution of oxalyl chloride (1.1 mL, 12.9 mmol) at -78 °C in dichloromethane (15 mL). The mixture was stirred for 15 min at -78 °C. A solution of 3-hydroxy-1,1-dimethylpropylcarbamic acid tert-butyl ester (1.75 g, 8.6 mmol) in dichloromethane (10 mL) was added dropwise over a period of 15 min. The solution was stirred at -78 °C for another 15 min. Triethylamine (6.0 mL, 43 mmol) was added. The solution was stirred at -78 °C for 5 min and then warmed to room temperature. The solution was diluted with dichloromethane (100 mL) and extracted with 1N hydrochloric acid (100 mL). The aqueous phase was extracted with dichloromethane (50 mL). The combined organic layers were washed with saturated sodium hydrogen carbonate solution (100 mL) and dried over magnesium sulfate. The solvent was removed in vacuo. The crude product was purified by column chromatography on silica (140 g) with ethyl acetate/heptane (1:3) to give 1.10 g of 3-(tert-butoxycarbonylamino)-3-methylbutanal.
- 25 MHz- $^1\text{H-NMR}$ (CDCl_3): d 1.39 (s, 6 H); 1.45 (s, 9 H); 2.85 (d, 2 H); 4.73 (br, 1 H); 9.80 (t, 1 H).

Ethyl (2E)-5-(tert-Butoxycarbonylamino)-5-methylhex-2-enoate:




- 5 Triethylphosphonoacetate (1.96 ml, 9.8 mmol) was dissolved in tetrahydrofuran (30 ml). Potassium tert-butoxide (1.10 g, 9.8 mmol) was added. The solution was stirred for 40 min at room temperature. A solution of 3-(tert-butoxycarbonylamino)-3-methylbutanal (1.10 g, 5.5 mmol) in Tetrahydrofuran (6 ml) was added. The solution was stirred at room temperature. for 75 min. It was diluted with ethyl acetate (100 ml) and 1N hydrochloric acid (100 ml). The
- 10 phases were separated. The aqueous phase was extracted with ethyl acetate (2 x 50 ml). The combined organic phases were washed with saturated sodium hydrogen carbonate solution (60 ml) and dried over magnesium sulfate. The solvent was removed in vacuo. The crude product was purified by column chromatography on silica (90 g) with ethyl acetate/heptane (1:4) to give 1.27 g of ethyl (2E)-5-(tert-butoxycarbonylamino)-5-methylhex-2-enoate.

15

¹H-NMR (CDCl₃): δ 1.30 (s, 6 H); 1.30 (t, 3 H); 1.46 (s, 9 H); 2.62 (d, 2 H); 4.27 (q, 2 H); 4.42 (br, 1 H); 5.88 (d, 1 H); 6.94 (td, 1 H).

- 20 (2E)-5-(tert-Butoxycarbonylamino)-5-methylhex-2-enoic acid:




- Ethyl (2E)-5-(tert-butoxycarbonylamino)-5-methylhex-2-enoate (1.233 g, 4.54 mmol) was
- 25 dissolved in dioxane (20 ml). Lithium hydroxide (0.120 g, 5.00 mmol) was added as a solid. Water (10 ml) was added, until a clear solution was reached. The solution was stirred 16 h at room temperature. The solution was diluted with water (70 ml) and was extracted with tert-butyl methyl ether (2 x 100 ml). The aqueous phase was acidified with 1N sodium hydrogensulfate

solution (pH = 1) and was extracted with tert-butylmethylether (3 x 70 ml). The organic phases were combined and dried over magnesium sulfate. The solvent was removed in vacuo to give 1.05 g of (2E)-5-(tert-butoxycarbonylamino)-5-methylhex-2-enoic acid. The crude product was used for further syntheses.


- 5 ¹H-NMR (DMSO d₆): d 1.15 (s, 6 H); 1.35 (s, 9 H); 2.53 (d, 2 H); 5.75 (d, 1 H); 6.57 (br, 1 H); 6.75 (td, 1 H); 12.15 (s, 1 H).

1-Aza-spiro[3.3]heptan-2-one:



- Methylenecyclobutane (40.0g, 0.587 mol) was dissolved in diethylether (250 ml). At - 40 °C chlorosulfonylisocyanate (26 ml, 0.294 mol) was added dropwise. The reaction mixture was warmed to 10 °C, An exothermic reaction was observed, and precipitation was formed. The reaction mixture was cooled to -20 °C. It was stirred for 16 h, while it was warming up to room temperature. A saturated aqueous solution of sodium sulfite (100 ml) was added dropwise. The reaction mixture was stirred vigorously for 1 h. Another saturated aqueous solution of sodium sulfite (100 ml) was added dropwise. Solid sodium hydrogen carbonate was added, until pH 7. Dichloromethane (500 ml) was added. The phases were separated. The organic layer was dried over magnesium sulfate. The solvent was removed in vacuo, to give 23.59 g of 1-aza-spiro[3.3]heptan-2-one.


¹H-NMR (CDCl₃): d 1.75 (m, 2 H); 2.26 (m, 2 H); 2.39 (m, 2 H); 2.96 (s, 2 H); 6.55 (br, 1 H).

2-Oxo-1-azaspiro[3.3]heptane-1-carboxylic acid *tert*-butylester

- 5 A solution of di-*tert*-butyl dicarbonate (55.7 g, 0.211 mol) in dichloromethane (100 ml) was added dropwise to a solution of 1-aza-spiro[3.3]heptan-2-one, triethylamine (36 ml, 0.255 mol), and 4-dimethylaminopyridine (2.6 g, 0.021 mol) in dichloromethane (100 ml). The reaction mixture was stirred for 16 h at room temperature. It was washed with a 10% aqueous solution of ammonium chloride (100 ml), water (100 ml) and a saturated aqueous solution of sodium hydrogen carbonate (100 ml). The organic layer was dried over magnesium sulfate.
- 10 The solvent was removed in vacuo to give 48.24 g of crude 2-oxo-1-azaspiro[3.3]heptane-1-carboxylic acid *tert*-butylester, which was used for the next step without purification.
- ¹H-NMR (CDCl₃): d 1.55 (s, 9 H); 1.78 (m, 1 H); 1.92 (m, 1 H); 2.18 (m, 2 H); 2.90 (m, 2 H); 3.04 (s, 1 H).

15

(1-(*tert*-Butoxycarbonylamino)cyclobutyl)acetic acid:




20

- An 1 N aqueous solution of lithium hydroxide (227 ml, 227 mmol) was added to a solution of 2-oxo-1-azaspiro[3.3]heptane-1-carboxylic acid *tert*-butylester (48 g, 0.227 mmol) in tetrahydrofuran (200 ml). The reaction mixture was stirred for 2 h. Diethyl ether (200 ml) and water (200 ml) were added. The mixture was stirred for 16 h. The organic layer was isolated. The
- 25 aqueous phase was extracted with diethyl ether (200 ml). The aqueous phase was acidified

with a 10% aqueous solution of sodium hydrogen sulfate until pH 3. The formed precipitation was filtered off, washed with water, and dried in vacuo, to give 38.84 g of (1-(*tert*-butoxycarbonylamino)cyclobutyl)acetic acid.

¹H-NMR (CDCl₃): d 1.45 (s, 9 H); 1.85 (m, 1 H); 1.95 (m, 1 H); 2.25 (m, 4 H); 2.87 (m, 2 H);
5 5.15 and 6.20 (both br, together 1 H).


(2*E*)-4-(1-(*tert*-Butoxycarbonylamino)cyclobutyl)but-2-enoic acid:



(2*E*)-4-(1-(*tert*-Butoxycarbonylamino)cyclobutyl)but-2-enoic acid was synthesized starting with (1-(*tert*-butoxycarbonylamino)cyclobutyl)acetic acid analogously to the synthesis of (2*E*)-5-(*tert*-butoxycarbonylamino)-5-methylhex-2-enoic acid starting with of 3-*tert*-butoxycarbonyl-
15 amino-3-methylbutanoic acid.

¹H-NMR (CDCl₃): d 1.43 (s, 9 H); 1.84 (m, 1 H); 1.95 (m, 1 H); 2.10 (m, 2 H); 2.20 (m, 2 H); 2.70 (m, 2 H); 4.75 (br, 0.5 H); 5.90 (m, 1 H); 6.35 (br, 0.5 H); 6.95 (m, 1 H).

(2*E*)-5-*tert*-Butoxycarbonylamino-3,5-dimethylhex-2-enoic acid ethyl ester:




Diacetonamine hydrogen oxalate (30.0 g; 146 mmol) was suspended in tetrahydrofuran (400
25 ml). An aqueous solution of sodium hydroxide (1 N; 146 ml) was added. Di-*tert*-butyl dicarbonate (38.3 g; 175 mmol) was dissolved in tetrahydrofuran (100 ml) and added dropwise to the reaction mixture. The reaction mixture was stirred for 2 h at room temperature. Sodium hydroxide (1 N; 146 ml) was added and the reaction mixture was stirred for 12 h at room

temperature. Water (200 ml) and ethyl acetate (200 ml) were added. The aqueous phase was extracted with ethyl acetate (4 x 200 ml). The combined organic phases were dried over magnesium sulfate, and the solvent was removed in vacuo. The residue was purified by flash chromatography on silica (200 g), using ethyl acetate/heptane (1:3) as eluent, to afford 28.4 g of (1,1-dimethyl-3-oxobutyl)carbamic acid *tert*-butyl ester.

Triethyl phosphonoacetate (4.7 g; 20.9 mmol) was dissolved in tetrahydrofuran (36 ml). Potassium *tert*-butoxide (2.3 g; 20.9 mmol) was added and the reaction mixture was stirred for 40 min at room temperature. (1,1-Dimethyl-3-oxobutyl)carbamic acid *tert*-butyl ester (2.5 g; 11.6 mmol) was dissolved in tetrahydrofuran (15 ml) and added dropwise to the reaction mixture which was heated to reflux for 12 h. Ethyl acetate (100 ml) and hydrochloric acid (1 N; 100 ml) were added and the phases were separated. The aqueous phase was extracted with ethyl acetate (3 x 50 ml). The combined organic phases were washed with an aqueous solution of sodium hydrogen carbonate (saturated; 100 ml), dried (magnesium sulfate) and evaporated in vacuo. The residue was purified by flash chromatography on silica (120 g) using ethyl acetate/heptane (1:2) as eluent to afford 2.0 g of (2E)-5-*tert*-butoxycarbonylamino-3,5-dimethylhex-2-enoic acid ethyl ester.

¹H-NMR (CDCl₃) δ 1.25 (t, 3H); 1.30 (s, 6H); 1.44 (s, 9H); 2.21 (s, 3H); 2.58 (s, 2H); 4.14 (q, 2H); 4.48 (s, 1H); 5.65 (s, 1H).

(2E)-5-*tert*-Butoxycarbonylamino-3,5-dimethylhex-2-enoic acid:




(2E)-5-*tert*-Butoxycarbonylamino-3,5-dimethylhex-2-enoic acid ethyl ester (1.95 g; 6.83 mmol) was dissolved in 1,4-dioxane (25 ml) and water (15 ml). Lithium hydroxide (0.18 g; 7.52 mmol) was added and the reaction mixture was stirred for 12 h at room temperature. Water (150 ml) and *tert*-butyl methyl ether (150 ml) was added. The aqueous phase was diluted with a 10% aqueous solution of sodium hydrogensulfate until pH 2.5 and extracted with *tert*-butyl methyl ether (3 x 100 ml). The combined organic phases were dried over magnesium sulfate and

evaporated in vacuo. The residue was recrystallized from heptane (20 ml) to afford 0.6 g of (2E)-5-*tert*-Butoxycarbonylamino-3,5-dimethylhex-2-enoic acid.

¹H-NMR (CDCl₃) δ 1.29 (s, 6H); 1.44 (s, 9H); 2.23 (s, 3H); 2.62 (s, 2H); 4.45 (s, 1H); 5.66 (s, 1H).

5

(2E)-5-(N-(*tert* Butoxycarbonyl)-N-methylamino)-5-methylhex-2-enoic acid.



10


(2E)-5-(*tert*-Butyloxycarbonylamino)-5-methylhex-2-enoic acid (5.00 g ; 20.6 mmol) was dissolved in tetrahydrofuran (70 ml). Methyl iodide (10.3 ml; 164 mmol) was added and the solution was cooled to 0° C. Sodium hydride (60% in oil) (2.07 g; 61.6 mmol) was added in portions and the solution was stirred at room temperature for four days. Ethyl acetate (70 ml) and water (60 ml) was added dropwise and the solvent was removed in vacuo. The crude product was dissolved in water (40 ml) and ether (40 ml). The organic phase was washed with a saturated aqueous solution of sodium hydrogencarbonate (30 ml). The aqueous phases were mixed and 5% aqueous citric acid was added to pH 3. The aqueous phase was extracted with ethylacetate (4 x 50 ml). The organic phase was washed with water (2 x 40 ml), an aqueous solution of sodium thiosulfate (5%; 40 ml), water (40 ml), dried over MgSO₄ and the solvent was removed in vacuo. The residue was dissolved in ethylacetate (45 ml) and washed with an aqueous solution of sodium hydrogensulfate (10%; 3 x 30 ml), dried over MgSO₄ and concentrated in vacuo to give 4.00 g of (2E)-5-(N-(*tert* Butoxycarbonyl)-N-methylamino)-5-methylhex-2-enoic acid.

25

¹H-NMR (CDCl₃) δ 1.38 (s, 6H), 1.45 (s, 9H); 2.80 (d, 2H); 2.85 (s, 3H); 5.88 (d, 1H); 7.01 (q, 1H).


Example 1

(2E)-5-Amino-5-methylhex-2-enoic acid N-((1R)-1-{N-[(1R)-1-benzyl-2-(4-
 ((dimethylamino)methyl)piperidin-1-yl)-2-oxoethyl]-N-methylcarbamoyl}-2-(2-naphthyl)ethyl)-
 5 N-methylamide



10

4-(Dimethylcarbamoyl)piperidine-1-carboxylic acid tert-butyl ester




15

1-(tert-Butoxycarbonyl)piperidine-4-carboxylic acid (8.0 g, 35 mmol) was dissolved in di-
 chloromethane (70 ml) and N,N-dimethylformamide (35 ml). 1-Hydroxy-7-azabenzotriazole
 (4.75 g, 35 mmol) was added. The solution was cooled to 0 °C. N-(3-dimethylaminopropyl)-
 N'-ethylcarbodiimide hydrochloride (6.69 g, 35 mmol) was added. The reaction mixture was
 20 stirred for 20 min at 0 °C. A 5.6 M solution of dimethylamine in ethanol (37 ml, 209 mmol)
 was added. The reaction mixture was stirred for 3 days, while it was warming up to room

temperature. It was diluted with ethyl acetate (400 ml) and washed with a 10% aqueous solution of sodium hydrogen sulfate (400 ml). The aqueous phase was extracted with ethyl acetate (2 x 200 ml). The combined organic layers were washed with a saturated aqueous solution of sodium hydrogen carbonate (300 ml) and dried over magnesium sulfate. The solvent was removed in vacuo. The crude product was purified by flash chromatography on silica (300 g), using dichloromethane/methanol 20:1 as eluent, to give 4.56 g of 4-(dimethylcarbamoyl)piperidine-1-carboxylic acid tert-butyl ester.

¹H-NMR (CDCl₃): δ 1.47 (s, 9 H); 1.70 (m, 4 H); 2.60 - 2.90 (m, 3 H); 2.96 (s, 3 H); 3.08 (s, 3 H); 4.17 (m, 2 H).


4-((Dimethylamino)methyl)piperidine-1-carboxylic acid tert-butyl ester



At 0 °C a solution of 4-((dimethylamino)methyl)piperidine-1-carboxylic acid tert-butyl ester (4.56 g, 18 mmol) in tetrahydrofuran (80 ml) was added dropwise to a suspension of sodium borohydride (1.61 g, 43 mmol) in tetrahydrofuran (80 ml). The reaction mixture was stirred for 20 min at 0 °C. A solution of iodine 4.51 g, 18 mmol) in tetrahydrofuran (80 ml) was added dropwise at 0 °C. The reaction mixture was heated to reflux for 16 h. It was cooled to 4 °C. Methanol (200 ml) was added dropwise. The solvent was removed in vacuo. The residue was dissolved in a 20% aqueous solution of sodium hydroxide (200 ml) and tert-butyl methyl ether (150 ml). The phases were separated. The aqueous phase was extracted with tert-butyl methyl ether (3 x 100 ml). The combined organic layers were dried over magnesium sulfate. The solvent was removed in vacuo. The crude product was purified by flash chromatography on silica (100 g), using dichloromethane/methanol/25% aqueous ammonia (100:10:1) as eluent, to give 4.07 g of 4-((dimethylamino)methyl)piperidine-1-carboxylic acid tert-butyl ester.

$^1\text{H-NMR}$ (CDCl_3): δ 1.22 (m, 2 H); 1.44 (s, 9 H); 1.85 (d, 2 H); 2.09 (m, 1 H); 2.61 (s, 6 H); 2.65 (m, 2 H); 2.78 (t, 2 H); 4.05 (d, 2 H).


5 N,N-Dimethyl-N-((piperidin-4-yl)methyl)amine



- 10 A 3 M solution of hydrogen chloride in ethyl acetate (120 ml, 360 mmol) was added to a solution of 4-((dimethylamino)methyl)piperidine-1-carboxylic acid tert-butyl ester (2.0 g, 14 mmol) in ethyl acetate (50 ml). The reaction mixture was stirred for 30 min at room temperature. The solvent was removed in vacuo to give 2.3 g of the crude dihydrochloride salt of N,N-dimethyl-N-((piperidin-4-yl)methyl)amine, which was used without purification for the
- 15 next step.

$^1\text{H-NMR}$ (CDCl_3 , selected values): δ 1.48 (m, 2 H); 1.92 (s, 6 H); 3.22 (d, 2 H).

- N-[(1R)-1-Benzyl-2-(4-((dimethylamino)methyl)piperidin-1-yl)-2-oxoethyl]-N-methylcarbamic acid tert-butyl ester
- 20




At 0 °C, N-(3-dimethylaminopropyl)-N'-ethylcarbodiimide hydrochloride (1.158 g, 6.04 mmol) was added to a solution of (2R)-2-(N-(tert-butoxycarbonyl)-N-methylamino)-3-phenylpropionic acid (1.69 g, 6.04 mmol) and 1-hydroxy-7-azabenzotriazole (0.822 g, 6.04 mmol) in dichloromethane (25 ml) and N,N-dimethylformamide (12 ml). The reaction mixture was stirred for 20 min at 0 °C. A solution of the crude dihydrochloride salt of N,N-dimethyl-N-((piperidin-4-yl)methyl)amine (1.3 g, 6.04 mmol) in N,N-dimethylformamide (10 ml) and dichloromethane (5 ml) and ethyldiisopropylamine (6.2 ml, 36.25 mmol) were added successively. The reaction mixture was stirred for 16 h, while it was warming up to room temperature. It was diluted with ethyl acetate (100 ml) and washed with a saturated aqueous solution of sodium hydrogen carbonate (100 ml). The organic layer was dried over magnesium sulfate. The solvent was removed in vacuo. The crude product was purified by flash chromatography on silica (100 g), using dichloromethane/methanol/25% aqueous ammonia (200:10:1) as eluent, to give 1.22 g of N-[(1R)-1-benzyl-2-(4-((dimethylamino)methyl)piperidin-1-yl)-2-oxoethyl]-N-methylcarbamic acid tert-butyl ester.

¹H-NMR (CDCl₃, selected values): δ 1.28, 1.11, 1.37, and 1.38 (all s, together 9 H); 4.00 (m, 1 H); 4.57 (m, 1 H); 4.97 and 5.28 (both t, together 1 H); 7.10 - 7.40 (m, 5 H).

MS: 404 [M+1]⁺

(2R)-1-(4-((Dimethylamino)methyl)piperidin-1-yl)-2-(methylamino)-3-phenylpropan-1-one




At 0 °C, trifluoroacetic acid (20 ml) was added to a solution of N-[(1R)-1-benzyl-2-(4-((dimethylamino)methyl)piperidin-1-yl)-2-oxoethyl]-N-methylcarbamic acid tert-butyl ester (1.22 g, 3.02 mmol) in dichloromethane (20 ml). The reaction mixture was stirred for 1 h at 0 °C.

The solvents were removed in vacuo. The residue was dissolved in dichloromethane (70 ml) and the solvent was removed in vacuo. The latter procedure was repeated two times. The crude product was purified by flash chromatography on silica (100 g), using dichloromethane/methanol/25% aqueous ammonia (100:10:1), to give 659 mg of (2R)-1-(4-
 5 ((dimethylamino)methyl)piperidin-1-yl)-2-(methylamino)-3-phenylpropan-1-one.

¹H-NMR (CDCl₃, selected values): δ 0.91 and 1.47 (m and d, together 1 H); 1.27 (m, 1 H); 4.62 (t, 1 H).

- 10 N-((1R)-1-{N-[(1R)-1-Benzyl-2-(4-((dimethylamino)methyl)piperidin-1-yl)-2-oxoethyl]-N-methylcarbamoyl}-2-(2-naphthyl)ethyl)-N-methylcarbamic acid tert-butyl ester




- 15 At 0 °C, N-(3-dimethylaminopropyl)-N'-ethylcarbodiimide hydrochloride (416 mg, 2.17 mmol) was added to a solution of (2R)-2-(N-(tert-butoxycarbonyl)-N-methylamino)-3-(2-naphthyl)-propioni acid (715 mg, 2.17 mmol) and 1-hydroxy-7-azabenzotriazole (296 mg, 2.17 mmol) in dichloromethane (20 ml) and N,N-dimethylformamide (10 ml). The reaction mixture was stirred for 20 min at 0 °C. A solution of (2R)-1-(4-((dimethylamino)methyl)piperidin-1-yl)-2-
 20 (methylamino)-3-phenylpropan-1-one (659 mg, 2.17 mmol) in dichloromethane (10 ml) and N,N-dimethylformamide (5 ml) and ethyldiisopropylamine (0.56 ml, 3.26 mmol) were added successively. The reaction mixture was stirred for 16 h, while it was warming up to room temperature. It was diluted with ethyl acetate (100 ml) and washed with a saturated aqueous solution of sodium hydrogen carbonate (100 ml). The aqueous solution was extracted with

ethyl acetate (3 x 50 ml). The combined organic layers were dried over magnesium sulfate. The solvent was removed in vacuo. The crude product was purified by flash chromatography on silica (80 g), using ethyl acetate/heptane/triethylamine (1:1:0.08) as eluent, to give 1.05 g of N-((1R)-1-{N-[(1R)-1-benzyl-2-(4-((dimethylamino)methyl)piperidin-1-yl)-2-oxoethyl]-N-methylcarbamoyl}-2-(2-naphthyl)ethyl)-N-methylcarbamic acid tert-butyl ester.

¹H-NMR (CDCl₃, selected values): δ 1.24 and 1.42 (both s, together 9 H); 5.04, 5.28, 5.44, 5.54, 5.73 (m, dd, dd, dd, and m, together 3 H);

(2R)-N-[(1R)-1-Benzyl-2-(4-((dimethylamino)methyl)piperidin-1-yl)-2-oxoethyl]-N-methyl-2-(methylamino)-3-(2-naphthyl)propionamide




At 0 °C, trifluoroacetic acid (18 ml) was added to a solution of N-((1R)-1-{N-[(1R)-1-benzyl-2-(4-((dimethylamino)methyl)piperidin-1-yl)-2-oxoethyl]-N-methylcarbamoyl}-2-(2-naphthyl)-ethyl)-N-methylcarbamic acid tert-butyl ester (1.05 g, 1.71 mmol) in dichloromethane (18 ml). The reaction mixture was stirred for 50 min at 0 °C. The solvents were removed in vacuo.

The residue was dissolved in dichloromethane (50 ml) and the solvent was removed in vacuo. The latter procedure was repeated two times. The crude product was purified by flash chromatography on silica (80 g), using dichloromethane/methanol/25% aqueous ammonia as eluent, to give 846 mg of (2R)-N-[(1R)-1-benzyl-2-(4-((dimethylamino)methyl)piperidin-1-yl)-2-oxoethyl]-N-methyl-2-(methylamino)-3-(2-naphthyl)propionamide.

¹H-NMR (CDCl₃, selected values): δ 0.60 (m, 1 H); 4.38 (t, 1 H); 5.72 and 5.79 (both t, together 1 H).

- 5 {(3E)-4-[N-((1R)-1-{N-[(1R)-1-Benzyl-2-(4-((dimethylamino)methyl)piperidin-1-yl)-2-oxoethyl]-N-methylcarbamoyl}-2-(2-naphthyl)ethyl)-N-methylcarbamoyl]-1,1-dimethylbut-3-enyl}carbamic acid tert-butyl ester



10

- At 0 °C, N-(3-dimethylaminopropyl)-N'-ethylcarbodiimide hydrochloride (112 mg, 0.58 mmol) was added to a solution of (2E)-5-(tert-butoxycarbonylamino)-5-methylhex-2-enoic acid (142 mg, 0.58 mmol) and 1-hydroxy-7-azabenzotriazole (79 mg, 0.58 mmol) in dichloromethane (10 ml) and N,N-dimethylformamide (5 ml). The reaction mixture was stirred for 20 min at 0 °C. A solution of (2R)-N-[(1R)-1-benzyl-2-(4-((dimethylamino)methyl)piperidin-1-yl)-2-oxoethyl]-N-methyl-2-(methylamino)-3-(2-naphthyl)propionamide (300 mg, 0.58 mmol) in dichloromethane (5 ml) and N,N-dimethylformamide (5 ml) and ethyldiisopropylamine 0.10 ml, 0.58 mmol) were added successively. The reaction mixture was stirred for 3 days, while it was warming up to room temperature. It was diluted with ethyl acetate (70 ml) and washed with a saturated aqueous solution of sodium hydrogen carbonate (70 ml). The aqueous phase was extracted with ethyl acetate (3 x 50 ml). The combined organic layers were dried over magnesium sulfate. The solvent was removed in vacuo. The crude product was purified by flash chromatography on silica (70 g), using dichloromethane/methanol/25% aqueous
- 15
- 20

ammonia (200:10:1) as eluent, to give 313 g of {(3E)-4-[N-((1R)-1-{N-[(1R)-1-benzyl-2-(4-((dimethylamino)methyl)piperidin-1-yl)-2-oxoethyl]-N-methylcarbamoyl}-2-(2-naphthyl)ethyl)-N-methylcarbamoyl]-1,1-dimethylbut-3-enyl}carbamic acid tert-butyl ester.

5 ¹H-NMR (CDCl₃, selected values): δ 1.28 and 1.30 (both s, together 6 H); 1.42 (s, 9H); 2.23, 2.27, 2.38, 2.43, 2.51, 2.52, 2.81, and 2.82 (all s, together 12 H); 5.56, 5.76, and 5.90 (m, m, and dd, together 2 H); 6.17 and 6.19 (both dd, together 1 H); 6.94 (m, 1 H).

10 At 0 °C, trifluoroacetic acid (6 ml) was added to a solution of {(3E)-4-[N-((1R)-1-{N-[(1R)-1-benzyl-2-(4-((dimethylamino)methyl)piperidin-1-yl)-2-oxoethyl]-N-methylcarbamoyl}-2-(2-naphthyl)ethyl)-N-methylcarbamoyl]-1,1-dimethylbut-3-enyl}carbamic acid tert-butyl ester (212 mg, 0.29 mmol) in dichloromethane (6 ml). The reaction mixture was stirred for 20 min at 0 °C. It was diluted with dichloromethane (30 ml). A saturated aqueous solution of sodium
15 hydrogen carbonate (30 ml) was added dropwise. Solid sodium hydrogen carbonate was added, until pH 7 was obtained. The phases were separated. The aqueous phase was extracted with dichloromethane (3 x 50 ml). The combined organic layers were dried over magnesium sulfate. The solvent was removed in vacuo. The crude product was purified by flash chromatography on silica (20 g), using dichloromethane/methanol/25% aqueous am-
20 monia (100:10:1) as eluent, to give 5 mg of the title compound.

¹H-NMR (CDCl₃, selected values): δ 1.20 (s, 6 H); 2.28, 2.32, 2.41, 2.49, 2.56, 2.57, 2.82, and 2.83 (all s, together 12 H); 5.58, 5.78, and, 5.92 (m, m, and dd, together 2 H); 6.16 and 6.19 (both d, together 1 H); 7.00 (m, 1 H).

25

HPLC: 39.23 min (A1).
41.55 min (B1).


MS: 640.4 [M+1]⁺.

30

For biological testing, the title compound was transferred into its acetate salt by lyophilization with 0.5 M acetic acid (40 ml).


Example 2

(2E)-5-Amino-5-methylhex-2-enoic acid N-((1R)-1-{N-[(1R)-1-benzyl-2-((3S)-3-(dimethylaminomethyl)piperidin-1-yl)-2-oxoethyl]-N-methylcarbamoyl}-2-(2-naphthyl)ethyl)-N-methylamide



10

(3R)-Piperidine-1,3-dicarboxylic acid 1-tert-butyl ester 3-ethyl ester




15

(R)-Ethyl nipetcotate tartrate (10.0 g, 32.5 mmol) were suspended in tetrahydrofuran (90 ml). An 1 N solution of sodium hydroxide in water (98 ml, 98 mmol) was added. A solution of di-tert-butyl dicarbonate (7.10 g, 32.5 mmol) in tetrahydrofuran (90 ml) was added. The reaction mixture was stirred for 16 h at room temperature. Ethyl acetate (400 ml) was added. The reaction mixture was washed with a 10% aqueous solution of sodium hydrogen sulfate (400

ml). The aqueous solution was extracted with ethyl acetate (2 x 200 ml). The combined organic layers were washed with a saturated aqueous solution of sodium hydrogen carbonate (200 ml) and dried over magnesium sulfate. The solvent was removed in vacuo. The crude product was purified by flash chromatography on silica (90 g), using ethyl acetate/heptane 1:4 as eluent, to give 4.13 g of (3R)-piperidine-1,3-dicarboxylic acid 1-tert-butyl ester 3-ethyl ester.

¹H-NMR (CDCl₃): δ 1.27 (t, 3 H); 1.48 (s, 9 H); 1.54 (m, 1 H); 1.62 (m, 1 H); 1.73 (m, 2H); 2.05 (m, 1 H); 2.45 (m, 1 H); 2.81 (m, 1H); 2.98 (br, 1 H); 3.93 (m, 1 H); 4.14 (q, 1 H).


(3R)-3-Formylpiperidine-1-carboxylic acid tert-butyl ester



A 1.2 M solution of diisobutylaluminum hydride in toluene (30.8 ml, 36.9 mmol) was added at -78 °C to a solution of (3R)-piperidine-1,3-dicarboxylic acid 1-tert-butyl ester 3-ethyl ester (4.13 g, 16.1 mmol) in diethyl ether (30 ml). The reaction mixture was stirred for 2.5 h at - 78 °C. Water (9.6 ml) was added dropwise. The reaction mixture was warmed to room temperature. The precipitation was removed by filtration through a plug of celite. The celite was washed with tert-butyl methyl ether (3 x 100 ml). The liquids were combined and dried over magnesium sulfate. The solvent was removed in vacuo, to give 1.94 g of crude (3R)-3-formylpiperidine-1-carboxylic acid tert-butyl ester, which was used for the next step without further purification.

¹H-NMR (CDCl₃): δ 1.45 (s, 9H); 1.67 (m, 2 H); 1.95 (m, 1 H); 2.43 (m, 1 H); 3.10 (m, 1 H); 3.32 (dd, 1 H); 3.52 (d, 1 H); 3.66 (m, 1 H); 3.95 (m, 1 H); 9.69 (s, 1 H).

5 (3S)-3-(Dimethylaminomethyl)piperidine-1-carboxylic acid tert-butyl ester




A solution of crude (3R)-3-formylpiperidine-1-carboxylic acid tert-butyl ester (1.94 g, 9.1 mmol) in dichloromethane (80 ml) was prepared. A 5.6 M solution of dimethylamine in ethanol (3.2 ml, 18.2 mmol) and molsieves were added successively. Sodium triacetoxyborohydride (5.78 g, 27.3 mmol) was added to this mixture. Acetic acid (1.04 ml, 18.2 mmol) was added. The reaction mixture was stirred for 16 h at room temperature. An 1 N aqueous solution of sodium hydroxide (70 ml) and tert-butyl methyl ether (70 ml) were added. The phases were separated. The aqueous solution was extracted with tert-butyl methyl ether (3 x 70 ml). The combined organic layers were dried over magnesium sulfate. The solvent was removed in vacuo. The crude product was purified by flash chromatography on silica (40 g), using dichloromethane/methanol/25% aqueous ammonia (100:10:1) as eluent, to give 866 mg of (3S)-3-(dimethylaminomethyl)piperidine-1-carboxylic acid tert-butyl ester.

20

¹H-NMR (CDCl₃): δ 1.10 (m, 1 H); 1.45 (s, 9 H); 1.45 (m, 1 H); 1.64 (m, 2 H); 1.85 (m, 1 H); 2.10 (m, 2 H); 2.20 (s, 6 H); 2.50 (br, 1 H); 2.79 (m, 1 H); 3.95 (m, 2 H).

N,N-Dimethyl-N-(((3R)-piperidin-3-yl)methyl)amine




- 5 (3S)-3-(Dimethylaminomethyl)piperidine-1-carboxylic acid tert-butyl ester (1.25 g, 5.15 mmol) was dissolved in ethyl acetate (30 ml). A 2.7 M solution of hydrogen chloride in ethyl acetate (75 ml, 203 mmol) was added. The reaction mixture was stirred for 45 min at room temperature. The solvent was removed in vacuo to give 976 mg of the crude dihydrochloride salt of N,N-dimethyl-N-(((3R)-piperidin-3-yl)methyl)amine, which was used for the next step
- 10 without further purification.

¹H-NMR (CD₃OD): δ 1.42 (m, 1 H); 1.86 (m, 1 H); 2.00 (m, 2 H); 2.38 (m, 1 H); 2.85 (t, 1 H); 2.95 (s, 6 H); 2.98 (m, 1 H); 3.16 (m, 2 H); 3.42 (m, 1 H); 3.53 (m, 1 H).

15

N-[(1R)-1-Benzyl-2-((3S)-3-(dimethylaminomethyl)piperidin-1-yl)-2-oxoethyl]-N-methylcarbamic acid tert-butyl ester



20

At 0 °C N-(3-dimethylaminopropyl)-N'-ethylcarbodiimide hydrochloride (870 mg, 4.54 mmol) was added to a solution (2R)-2-(N-(tert-butoxycarbonyl)-N-methylamino)-3-phenylpropionic acid (1.27 g, 4.54 mmol) and 1-hydroxy-7-azabenzotriazole (617 mg, 4.54 mmol) in dichloro-

methane (20 ml) and N,N-dimethylformamide (10 ml). The reaction mixture was stirred for 20 min at 0 °C. A solution of the crude dihydrochloride salt of N,N-dimethyl-N-(((3R)-piperidin-3-yl)methyl)amine (976 mg, 4.54 mmol) in dichloromethane (20 ml) and N,N-dimethylformamide (10 ml) and ethyldiisopropylamine (3.9 ml, 22.7 mmol) were added successively.

5 The reaction mixture was stirred for 3 days, while it was warming up to room temperature. Ethyl acetate (300 ml) was added. The solution was washed with a saturated aqueous solution of sodium hydrogen carbonate (300 ml). The aqueous phase was extracted with ethyl acetate (2 x 200 ml). The combined organic layers were dried over magnesium sulfate. The solvent was removed in vacuo. The crude product was purified by flash chromatography on


10 silica (90 g), using dichloromethane/methanol/25% aqueous ammonia (100:10:1) as eluent, to give 1.69 g of N-[(1R)-1-benzyl-2-((3S)-3-(dimethylaminomethyl)piperidin-1-yl)-2-oxoethyl]-N-methylcarbamic acid tert-butyl ester.

¹H-NMR (CDCl₃, selected values): δ 1.20, 1.24, 1.31, and 1.32 (all s, together 9 H); 2.12, 2.13, and 2.18 (all s, together 6 H); 2.81 (m, 3 H); 4.97 and 5.30 (both m, together 1 H); 7.05 - 7.35 (m, 5 H).

15

(2R)-1-((3S)-3-((Dimethylamino)methyl)piperidin-1-yl)-2-methylamino-3-phenylpropan-1-one

20



At 0 °C, trifluoroacetic acid (25 ml) was added to a solution of N-[(1R)-1-benzyl-2-((3S)-3-(dimethylaminomethyl)piperidin-1-yl)-2-oxoethyl]-N-methylcarbamic acid tert-butyl ester (1.69 g, 4.2 mmol) in dichloromethane (25 ml). The reaction mixture was stirred for 30 min at 0 °C. The solvent was removed in vacuo. The residue was dissolved in dichloromethane (100 ml)

25


and the solvent was removed in vacuo. The latter procedure was repeated two times. The crude product was purified by flash chromatography on silica (90 g), using dichloromethane/methanol/25% aqueous ammonia (100:10:1) as eluent, to give 1.15 g of (2R)-1-((3S)-3-((dimethylamino)methyl)piperidin-1-yl)-2-methylamino-3-phenylpropan-1-one.

5

¹H-NMR (CDCl₃, selected values): δ 0.38, 1.11, 1.37, and 1.65 (all m, together 4 H); 2.11, 2.19, 2.25, and 2.31 (all s, together 9 H); 4.37 and 4.53 (both m, together 1 H); 7.10 - 7.35 (m, 5 H).

10

N-((1R)-1-{N-[(1R)-1-Benzyl-2-((3S)-3-((dimethylamino)methyl)piperidin-1-yl)-2-oxoethyl]-N-methylcarbamoyl}-2-(2-naphthyl)ethyl)-N-methylcarbamic acid tert-butyl ester



15


At 0 °C N-(3-dimethylaminopropyl)-N'-ethylcarbodiimide hydrochloride (379 mg, 1.98 mmol) was added to a solution of (2R)-2-(N-(tert-butoxycarbonyl)-N-methylamino)-3-(2-naphthyl)propionic acid (651 mg, 1.98 mmol) and 1-hydroxy-7-azabenzotriazole (269 mg, 1.98 mmol) in dichloromethane (10 ml) and N,N-dimethylformamide (5 ml). The reaction mixture was stirred for 20 min at 0 °C. A solution of (2R)-1-((3S)-3-((dimethylamino)methyl)piperidin-1-yl)-2-methylamino-3-phenylpropan-1-one (600 mg, 1.98 mmol) in dichloromethane (10 ml) and ethyldiisopropylamine (0.51 ml, 2.97 mmol) were added successively. The reaction mixture was stirred for 3 days, while it was warming up to room temperature. Ethyl acetate (100 ml)

20

was added. The solution was washed with a saturated aqueous solution of sodium hydrogen carbonate (100 ml). The aqueous phase was extracted with ethyl acetate (3 x 50 ml). The combined organic layers were dried over magnesium sulfate. The solvent was removed in vacuo. The crude product was purified by flash chromatography on silica (90 g), using dichloromethane/methanol/25% aqueous ammonia (100:10:1) as eluent, to give 1.18 g of N-((1R)-1-{N-[(1R)-1-benzyl-2-((3S)-3-((dimethylamino)methyl)piperidin-1-yl)-2-oxoethyl]-N-methylcarbamoyl}-2-(2-naphthyl)ethyl)-N-methylcarbamamic acid tert-butyl ester.

¹H-NMR (CDCl₃, selected values): δ 0.45 and 0.71 (both m, together 1 H); 1.03, 1.05, 1.15, 1.20, 1.28, 1.36, and 1.42 (all s, together 9 H); 2.12, 2.15, 2.21, 2.26, 2.29, 2.85 (all s, together 6 H); 5.05, 5.44, 5.58, 5.71, 5.85, and 6.00 (all s, together 2 H); 7.10 - 7.80 (m, 12 H).

(2R)-N-[(1R)-1-Benzyl-2-((3S)-3-((dimethylamino)methyl)piperidin-1-yl)-2-oxoethyl]-N-methyl-2-(methylamino)-3-(2-naphthyl)propionamide




At 0 °C, trifluoroacetic acid (20 ml) was added to a solution of N-((1R)-1-{N-[(1R)-1-benzyl-2-((3S)-3-((dimethylamino)methyl)piperidin-1-yl)-2-oxoethyl]-N-methylcarbamoyl}-2-(2-naphthyl)ethyl)-N-methylcarbamamic acid tert-butyl ester (1.18g, 1.92 mmol) in dichloromethane (20 ml). The reaction mixture was stirred for 50 min at 0 °C. The solvent was removed in vacuo. The residue was dissolved in dichloromethane (80 ml) and the solvent was

removed in vacuo. The latter procedure was repeated two times. The crude product was purified by flash chromatography on silica (40 g), using dichloromethane/methanol/25% aqueous ammonia (100:10:1) as eluent, to give 788 mg of (2R)-N-[(1R)-1-benzyl-2-((3S)-3-((dimethylamino)methyl)piperidin-1-yl)-2-oxoethyl]-N-methyl-2-(methylamino)-3-(2-naphthyl)-propionamide.

¹H-NMR (CDCl₃, selected values): δ 2.01 and 2.25 (both s, together 9 H); 3.72 (m, 2 H); 3.95 and 4.27 (both m, together 1 H); 5.77, 5.86, and 6.03 (t, m, and dd, together 1 H); 7.10 and 7.85 (m, 12 H).

{(3E)-4-[N-((1R)-1-{N-[(1R)-1-Benzyl-2-((3S)-3-((dimethylamino)methyl)piperidin-1-yl)-2-oxoethyl]-N-methylcarbamoyl}-2-(2-naphthyl)ethyl)-N-methylcarbamoyl]-1,1-dimethylbut-3-enyl}carbamic acid tert-butyl ester



At 0 °C, N-(3-dimethylaminopropyl)-N'-ethylcarbodiimide hydrochloride (105 mg, 0.55 mol) was added to a solution of (2E)-5-(tert-butoxycarbonylamino)-5-methylhex-2-enoic (136 mg, 0.55 mmol) and 1-hydroxy-7-azabenzotriazole (74 mg, 0.55 mmol) in dichloromethane (5 ml). The reaction mixture was stirred for 20 min at 0 °C. A solution of (2R)-N-[(1R)-1-benzyl-2-((3S)-3-((dimethylamino)methyl)piperidin-1-yl)-2-oxoethyl]-N-methyl-2-(methylamino)-3-(2-naphthyl)propionamide (281 mg, 0.55 mmol) in dichloromethane (5 ml) and N,N-dimethylformamide (5 ml) and ethyldiisopropylamine (0.094 ml, 0.55 mmol) were added suc-

cessively. The reaction mixture was stirred for 16 h, while it was warming up to room temperature. It was diluted with ethyl acetate (70 ml) and washed with a saturated aqueous solution of sodium hydrogen carbonate (70 ml). The aqueous phase was extracted with ethyl acetate (3 x 50 ml). The combined organic layers were dried over magnesium sulfate. The solvent was removed in vacuo. The crude product was purified by flash chromatography on silica (40 g), using dichloromethane/methanol/25% aqueous ammonia (100:10:1) as eluent, to give 398 mg of {(3E)-4-[N-((1R)-1-{N-[(1R)-1-benzyl-2-((3S)-3-((dimethylamino)methyl)piperidin-1-yl)-2-oxoethyl]-N-methylcarbamoyl]-2-(2-naphthyl)ethyl)-N-methylcarbamoyl]-1,1-dimethylbut-3-enyl}carbamic acid tert-butyl ester.

¹H-NMR (CDCl₃, selected values): δ 1.44 (s, 9 H); 5.58, 5.75, and 5.86 (all m, 2 H); 6.09 and 6.17 (both d, together 1 H); 6.84 (m, 1 H); 7.10 - 7.80 (m, 12 H).

At 0 °C, trifluoroacetic acid (7 ml) was added to a solution of {(3E)-4-[N-((1R)-1-{N-[(1R)-1-benzyl-2-((3S)-3-((dimethylamino)methyl)piperidin-1-yl)-2-oxoethyl]-N-methylcarbamoyl]-2-(2-naphthyl)ethyl)-N-methylcarbamoyl]-1,1-dimethylbut-3-enyl}carbamic acid tert-butyl ester (398 mg, 0.54 mmol) in dichloromethane (7 ml). The reaction mixture was stirred for 40 min at 0 °C. The solvent was removed in vacuo. The residue was dissolved in dichloromethane (20 ml) and the solvent was removed in vacuo. The latter procedure was repeated two times. The crude product was purified by flash chromatography on silica (40 g), using dichloromethane/methanol/25% aqueous ammonia (100:10:1) as eluent, to give 150 mg of the title compound.

¹H-NMR (CDCl₃, selected values): δ 1.08, 1.12, 1.14, and 1.15 (all s, together 6 H), 5.46, 5.59, 5.75, and 5.94 (all m, together 2 H); 6.15 (m, 1 H); 6.93 (m, 1 H).

HPLC 27.55 min (A1).
30.23min (B1).


LC-MS: 640.4 [M+1]⁺ at 8.54 min.

For biological testing, the title compound was transferred into its acetate salt, by lyophilization from 0.5 M acetic acid (40 ml).


5 Example 3

(2E)-4-(1-Aminocyclobutyl)but-2-enoic acid N-((1R)-1-{N-[(1R)-1-benzyl-2-((3S)-3-(dimethylaminomethyl)piperidin-1-yl)-2-oxoethyl]-N-methylcarbamoyl}-2-(2-naphthyl)ethyl)-N-methylamide

10



(1-((2E)-3-[N-((1R)-1-{N-[(1R)-1-Benzyl-2-((3S)-3-((dimethylamino)methyl)piperidin-1-yl)-2-oxoethyl]-N-methylcarbamoyl}-2-(2-naphthyl)ethyl)-N-methylcarbamoyl]allyl)cyclobutyl)carbamic acid tert-butyl ester



5

At 0 °C, N-(3-dimethylaminopropyl)-N'-ethylcarbodiimide hydrochloride (85 mg, 0.44 mmol) was added to a solution of (2E)-5-(tert-butoxycarbonylamino)-5-methylhex-2-enoic acid (113 mg, 0.44 mmol) and 1-hydroxy-7-azabenzotriazole (60 mg, 0.44 mmol) in dichloromethane (5 ml) and N,N-dimethylformamide (5 ml). The reaction mixture was stirred for 20 min at 0 °C. A solution of (2R)-N-[(1R)-1-benzyl-2-((3S)-3-((dimethylamino)methyl)piperidin-1-yl)-2-oxoethyl]-N-methyl-2-(methylamino)-3-(2-naphthyl)propionamide (228 mg, 0.44 mmol) in dichloromethane (10 ml) and ethyldiisopropylamine (0.07 ml, 0.44 mmol) were added successively. The reaction mixture was stirred for 16 h, while it was warming up to room temperature. It was diluted with ethyl acetate (70 ml) and washed with a saturated aqueous solution of sodium hydrogen carbonate (70 ml). The aqueous phase was extracted with ethyl acetate (3 x 50 ml). The combined organic layers were dried over magnesium sulfate. The solvent was removed in vacuo. The crude product was purified by flash chromatography on silica (40 g), using dichloromethane/methanol/25% aqueous ammonia (100:10:1) as eluent, to give 314 mg of (1-((2E)-3-[N-((1R)-1-{N-[(1R)-1-benzyl-2-((3S)-3-((dimethylamino)methyl)piperidin-1-yl)-2-oxoethyl]-N-methylcarbamoyl}-2-(2-naphthyl)ethyl)-N-methylcarbamoyl]allyl)cyclobutyl)carbamic acid tert-butyl ester.

20

¹H-NMR (CDCl₃, selected values): δ 1.40 (m, 9 H); 4.22 and 4.67 (both m, together 2 H); 5.60, 5.75, 5.85, and 5.90 (dd, dd, m, and m, together 2 H); 6.10 and 6.19 (both d, together 1 H); 6.73 and 6.87 (both m, together 1 H); 7.22, 7.42, and 7.76 (all m, together 12 H).

5

At 0 °C, trifluoroacetic acid (7 ml) was added to a solution of (1-((2E)-3-[N-((1R)-1-{N-[(1R)-1-benzyl-2-((3S)-3-((dimethylamino)methyl)piperidin-1-yl)-2-oxoethyl]-N-methylcarbamoyl]-2-(2-naphthyl)ethyl)-N-methylcarbamoyl]allyl)cyclobutyl)carbamic acid tert-butyl ester (314 mg, 0.42 mmol) in dichloromethane (7 ml). The reaction mixture was stirred for 20 min at 0 °C.

10

The solvent was removed in vacuo without heating. The residue was dissolved in dichloromethane (20 ml) and the solvent was removed in vacuo. The latter procedure was repeated two times. The crude product was purified by flash chromatography on silica (40 g), using dichloromethane/methanol/ammonia (100:10:1) as eluent, to give 180 mg of the title compound.

15

¹H-NMR (CDCl₃, selected values): δ 0.40 and 0.74 (both m, together 2 H); 3.73 and 4.22 (both m, together 2 H); 5.57, 5.77, and 5.91 (all m, together 2 H); 6.15 and 6.24 (both d, together 1 H); 6.85 and 6.96 (both m, together 1 H); 7.22, 7.92, and 7.74 (all m, together 12 H).

20

HPLC: 28.03 min (A1).
29.92min (B1).

MS: 652.4 [M+1]⁺


25

For biological testing, the title compound was transferred into its diacetate salt, by lyophilization from 0.5 M acetic acid (40 ml).

30


Example 4

(2E)-5-Amino-5-methylhex-2-enoic acid N-((1R)-1-{N-[(1R)-1-benzyl-2-((2S)-2-
 ((dimethylamino)methyl)pyrrolidin-1-yl)-2-oxoethyl]-N-methylcarbamoyl}-2-(2-naphthyl)ethyl)-
 5 N-methylamide



10

(2S)-Pyrrolidine-1,2-dicarboxylic acid 1-tert-butyl ester 2-ethyl ester



15


N-tert-Butoxycarbonylprolin (24.38 g, 113 mmol) was dissolved in dichloromethane (60 ml). Ethanol (7.9 ml, 135 mmol) and 4-dimethylaminopyridine (1.52 g, 12.5 mmol) were added. The solution was cooled to 0 °C. N-(3-dimethylaminopropyl)-N'-ethylcarbodiimide hydrochloride (23.88 g, 125 mmol) was added. The reaction mixture was stirred for 16 h, while it was

20

warming up to room temperature. Ethyl acetate (400 ml) was added. It was washed with a 10% aqueous solution of sodium hydrogen sulfate (300 ml). The aqueous phase was extracted with ethyl acetate (3 x 200 ml). The combined organic layers were washed with a saturated aqueous solution of sodium hydrogen carbonate (300 ml) and dried over magnesium sulfate. The solvent was removed in vacuo. The crude product was purified by flash chromatography on silica (400 g), using ethyl acetate (1:4) as eluent, to give 17.11 g of (2S)-pyrrolidine-1,2-dicarboxylic acid 1-tert-butyl ester 2-ethyl ester.

¹H-NMR (CDCl₃): δ 1.28 (m, 3 H); 1.43 and 1.46 (both s, together 9 H); 2.95 (m, 3 H); 2.22 (m, 1 H); 3.50 (m, 2 H); 4.18 and 4.30 (m and dd, together 3 H).


N-t-Butyloxycarbonyl-(S)-prolinal



At -78 °C, a 1.2 M solution of diisobutylaluminum hydride (31.7 ml, 38 mmol) in toluene was added dropwise to a solution of (2S)-pyrrolidine-1,2-dicarboxylic acid 1-tert-butyl ester 2-ethyl ester (4.02 g, 16.5 mmol) in diethyl ether (15 ml). The reaction mixture was stirred for 3 h at -78 °C. Water (9.9 ml) was added dropwise. The reaction mixture was warmed to room temperature. The mixture was filtered through a plug of celite. The celite was washed with tert-butyl methyl ether (3 x 100 ml). The combined organic layers were dried over magnesium sulfate. The solvent was removed in vacuo, to give 2.34 g of crude N-t-butyloxy-carbonyl-(S)-prolinal, which was used for the next step without further purification.

¹H-NMR (CDCl₃): δ 1.42 and 1.47 (both s, together 9 H); 1.70 - 2.20 (m, 4 H); 3.20 - 4.30 (m, 3 H); 9.45 and 9.55 (both s, together 1 H).

(2S)-2-((Dimethylamino)methyl)pyrrolidine-1-carboxylic acid tert-butyl ester



5

Crude N-t-butyloxycarbonyl-(S)-prolinal (2.34 g, 11.7 mmol) was dissolved in dichloromethane (90 ml). A 5.6 M solution of dimethylamine in ethanol (4.19 ml, 23.5 mmol) was added. 0.4 nm Mol sieves (10.0g) was added. Sodium triacetoxyborohydride 7.47 g, 35.2 mmol) and glacial acetic acid (1.34 ml, 23.5 mmol) were added successively. The reaction mixture was stirred for 3 days. It was filtered through a plug of celite. The celite was washed with methanol (150 ml). An 1N aqueous solution of sodium hydroxide (150 ml) and tert-butyl methyl ether (150 ml) were added. The phases were separated. The aqueous phase was extracted with tert-butyl methyl ether (3 x 100 ml). The combined organic layers were dried over magnesium sulfate. The solvent was removed in vacuo. The crude product was purified by flash chromatography on silica (90 g), using dichloromethane/methanol/25% aqueous ammonia (100:10:1) as eluent, to give 1.29 g of (2S)-2-((dimethylamino)methyl)pyrrolidine-1-carboxylic acid tert-butyl ester.


10

15

20

¹H-NMR (CDCl₃): δ 1.48 (s, 9 H); 1.90 (m, 4 H); 2.15 and 2.23 (AB, 2 H); 2.26 (s, 6 H); 3.31 (br, 2 H); 3.85 (br, 1 H).

N-Dimethyl-N-(((2S)-pyrrolidin-2-yl)methyl)amine




25

A 2.7 M solution of hydrogen chloride in ethyl acetate (75 ml, 202 mmol) was given to a solution of (2S)-2-((dimethylamino)methyl)pyrrolidine-1-carboxylic acid tert-butyl ester (1.29 g, 5.65 mmol) in ethyl acetate (30 ml). The reaction mixture was stirred for 30 min at room temperature. The solvent was removed in vacuo to give 1.36 g of the crude dihydrochloride salt of N-dimethyl-N-(((2S)-pyrrolidin-2-yl)methyl)amine, which was used for the next step without further purification.

¹H-NMR (CDCl₃): δ 1.90 (m, 2 H); 2.17 (m, 1 H); 2.40 (m, 1 H); 2.90 (m, 2 H); 3.14 (s, 6 H); 3.55 (m, 2 H); 4.35 (m, 1 H).

N-[(1R)-1-Benzyl-2-((2S)-2-((dimethylamino)methyl)pyrrolidin-1-yl)-2-oxoethyl]-N-methylcarbamic acid tert-butyl ester



At 0 °C, N-(3-dimethylaminopropyl)-N'-ethylcarbodiimide hydrochloride (1.30 g, 6.76 mmol) was added to a solution of (2R)-2-(N-(tert-butoxycarbonyl)-N-methylamino)-3-phenylpropionic acid (1.89 g, 6.76 mmol) and 1-hydroxy-7-azabenzotriazole 0.92 g, 6.76 mmol) in dichloromethane (10 ml). The reaction mixture was stirred for 20 min at 0 °C. A solution of the crude dihydrochloride salt of N-dimethyl-N-(((2S)-pyrrolidin-2-yl)methyl)amine (1.36 g, 6.76 mmol) in dichloromethane (10 ml) and N,N-dimethylformamide (10 ml) and ethyldiisopropylamine (5.75 ml, 33.8 mmol) were added successively. The reaction mixture was stirred for 16 h, while it was warming up to room temperature. It was diluted with ethyl acetate (100 ml) and washed with a saturated aqueous solution of sodium hydrogen carbonate (100 ml) and washed with a saturated aqueous solution of sodium hydrogen carbonate (100 ml). The aqueous phase was extracted with ethyl acetate (3 x 80 ml). The combined organic layers were dried over magnesium sulfate. The solvent was removed in vacuo. The crude product was purified by flash chromatography on silica (90 g), using dichloromethane/methanol/25% aqueous ammonia (100:10:1) as eluent, to give 2.26 g of N-[(1R)-1-

benzyl-2-((2S)-2-((dimethylamino)methyl)pyrrolidin-1-yl)-2-oxoethyl]-N-methylcarbamic acid tert-butyl ester.

¹H-NMR (CDCl₃, selected values): δ 1.20, 1.33, and 1.37 (all s, together 9 H); 2.22 and 2.28 (both s, together 6 H); 2.82 and 2.84 (both s, together 3 H); 4.25 (m, 1 H); 4.80, 5.11, and 5.30 (dd, t, and m, together 1 H); 7.10 - 7.30 (m, 5 H).

C₂₂H₃₅N₃O₃

[389.5]


calc.

C67.83 H9.06 N10.79

found

C67.39 H9.13 N10.73

(2R)-1-((2S)-2-((Dimethylamino)methyl)pyrrolidin-1-yl)-2-methylamino-3-phenylpropan-1-one




At 0 °C, trifluoroacetic acid (8 ml) was added to a solution of N-[(1R)-1-benzyl-2-((2S)-2-((dimethylamino)methyl)pyrrolidin-1-yl)-2-oxoethyl]-N-methylcarbamic acid tert-butyl ester (2.26 g, 5.80 mmol) in dichloromethane (8 ml). The reaction mixture was stirred for 20 min at 0 °C. The solvent was removed in vacuo. Dichloromethane (70 ml) was added, and the solvent was removed in vacuo. The latter procedure was repeated two times. The crude product was purified by flash chromatography on silica (90 g), using dichloromethane/methanol/25% aqueous ammonia (100:10:1) as eluent, to give 1.24 g of (2R)-1-((2S)-2-((dimethylamino)methyl)pyrrolidin-1-yl)-2-methylamino-3-phenylpropan-1-one.

¹H-NMR (CDCl₃, selected values): δ 2.33 (s, 3 H); 2.43 (s, 6 H); 3.25 (m, 3 H); 4.17 (m, 1 H); 7.25 (m, 5 H).

N-((1R)-1-{N-[(1R)-1-Benzyl-2-((2S)-2-((dimethylamino)methyl)pyrrolidin-1-yl)-2-oxoethyl]-N-methylcarbamoyl}-2-(2-naphthyl)ethyl)-N-methylcarbamic acid tert-butyl ester

5




At 0 °C, N-(3-dimethylaminopropyl)-N'-ethylcarbodiimide hydrochloride (530 mg, 2.76 mmol) was added to a solution of (2R)-2-(N-(tert-butoxycarbonyl)-N-methylamino)-3-(2-naphthyl)-propionic acid (911 mg, 2.76 mmol) and 1-hydroxy-7-azabenzotriazole (376 mg, 2.76 mmol) in dichloromethane (5 ml). The reaction mixture was stirred for 20 min at 0 °C. A solution of (2R)-1-((2S)-2-((dimethylamino)methyl)pyrrolidin-1-yl)-2-methylamino-3-phenylpropan-1-one (800 mg, 2.76 mmol) in dichloromethane (5 ml) and N,N-dimethylformamide (5 ml) and ethyldiisopropylamine (0.71 ml, 4.15 mmol) were added successively. The reaction mixture was stirred for 3 days, while it was warming up to room temperature. It was diluted with ethyl acetate (100 ml) and washed with a saturated aqueous solution of sodium hydrogen carbonate (100 ml). The aqueous solution was extracted with ethyl acetate (3 x 70 ml). The combined organic layers were dried over magnesium sulfate. The solvent was removed in vacuo. The crude product was purified by flash chromatography on silica (90 g), using dichloromethane/methanol/25% aqueous ammonia (200:10:1) as eluent, to give 1.37 g of N-((1R)-1-{N-[(1R)-1-benzyl-2-((2S)-2-((dimethylamino)methyl)pyrrolidin-1-yl)-2-oxoethyl]-N-methylcarbamoyl}-2-(2-naphthyl)ethyl)-N-methylcarbamic acid tert-butyl ester.

¹H-NMR (CDCl₃, selected values): δ 0.64 (m, 1 H); 1.10, 1.29, 1.36, and 1.47 (all s, together 9 H); 4.99, 5.09, 5.45, and 5.53 (t, t, m, and t, together 2 H); 7.10 - 7.90 (m, 12 H).

(2R)-N-[(1R)-1-Benzyl-2-((2S)-2-((dimethylamino)methyl)pyrrolidin-1-yl)-2-oxoethyl]-N-methyl-2-(methylamino)-3-(2-naphthyl)propionamide

5




At 0 °C, trifluoroacetic acid (10 ml) was added to a solution of N-((1R)-1-{N-[(1R)-1-benzyl-2-((2S)-2-((dimethylamino)methyl)pyrrolidin-1-yl)-2-oxoethyl]-N-methylcarbamoyl}-2-(2-naphthyl)ethyl)-N-methylcarbamic acid tert-butyl ester (1.37 g, 2.28 mmol) in dichloromethane (10 ml). The reaction mixture was stirred for 75 min at 0 °C. The solvent was removed in vacuo. The residue was dissolved in dichloromethane (70 ml) and the solvent was removed in vacuo. The latter procedure was repeated two times. The crude product was purified by flash chromatography on silica (90 g), using dichloromethane/methanol/25% aqueous ammonia (100:10:1) as eluent, to give 692 mg of (2R)-N-[(1R)-1-benzyl-2-((2S)-2-((dimethylamino)methyl)pyrrolidin-1-yl)-2-oxoethyl]-N-methyl-2-(methylamino)-3-(2-naphthyl)propionamide.

¹H-NMR (CDCl₃, selected values): δ 1.85 and 2.01 (both s, together 3 H); 2.20 and 2.31

(both s, together 6 H); 3.65 and 3.80 (both t, 1 H); 4.04 and 4.45 (both m, together 1 H); 5.60 and 5.91 (t and dd, together 1 H); 7.10 - 7.90 (m, 12 H).

{{(3E)-4-[N-((1R)-1-[N-[(1R)-1-Benzyl-2-((2S)-2-((dimethylamino)methyl)pyrrolidin-1-yl)-2-oxoethyl]-N-methylcarbamoyl]-2-(2-naphthyl)ethyl)-N-methylcarbamoyl]-1,1-dimethylbut-3-enyl}carbamic acid tert-butyl ester



5

At 0°C, N-(3-dimethylaminopropyl)-N'-ethylcarbodiimide hydrochloride (132 mg, 0.69 mmol) was added to a solution of (2E)-5-(tert-butoxycarbonylamino)-5-methylhex-2-enoic acid (168 mg, 0.69 mmol) and 1-hydroxy-7-azabenzotriazole (94 mg, 0.69 mmol) in dichloromethane (5 ml). The reaction mixture was stirred for 20 min at 0 °C. A solution of (2R)-N-[(1R)-1-benzyl-2-((2S)-2-((dimethylamino)methyl)pyrrolidin-1-yl)-2-oxoethyl]-N-methyl-2-(methylamino)-3-(2-naphthyl)propionamide (345 mg, 0.69 mmol) in dichloromethane (10 ml) and N,N-dimethylformamide (5 ml) and ethyldiisopropylamine were added successively. The reaction mixture was stirred for 16 h, while it was warming up to room temperature. It was diluted with ethyl acetate (70 ml) and washed with a saturated aqueous solution of sodium hydrogen carbonate (70 ml). The aqueous phase was extracted with ethyl acetate (3 x 50 ml). The combined organic layers were dried over magnesium sulfate. The solvent was removed in vacuo. The crude product was purified by flash chromatography on silica (40 g), using dichloromethane/methanol/25% aqueous ammonia (100:10:1) as eluent, to give 491 mg of

{{(3E)-4-[N-((1R)-1-[N-[(1R)-1-benzyl-2-((2S)-2-((dimethylamino)methyl)pyrrolidin-1-yl)-2-oxoethyl]-N-methylcarbamoyl]-2-(2-naphthyl)ethyl)-N-methylcarbamoyl]-1,1-dimethylbut-3-enyl}carbamic acid tert-butyl ester.

20

¹H-NMR (CDCl₃, selected values): δ 1.30 and 1.32 (both s, together 6 H); 1.45 (s, 9 H); 1.60 (m, 2 H); 4.00 (m, 1 H); 4.48 (m, 1 H); 5.48 (dd, 1 H); 5.92 (dd, 1 H); 6.11 and 6.20 (both d, together 1 H); 6.82 and 6.92 (both m, together 1 H); 7.10 - 7.90 (m, 12 H).

5 At 0 °C, trifluoroacetic acid (7 ml) was added to a solution of {(3E)-4-[N-((1R)-1-{N-[(1R)-1-benzyl-2-((2S)-2-((dimethylamino)methyl)pyrrolidin-1-yl)-2-oxoethyl]-N-methylcarbamoyl}-2-(2-naphthyl)ethyl)-N-methylcarbamoyl]-1,1-dimethylbut-3-enyl}carbamic acid tert-butyl ester (491 mg, 0.68 mmol) in dichloromethane (7 ml). The reaction mixture was stirred for 60 min at 0 °C. The solvent was removed in vacuo. The residue was dissolved in dichloromethane
10 (100 ml) and the solvent was removed in vacuo. The latter procedure was repeated two times. The crude product was purified by flash chromatography on silica (40 g), using dichloromethane/methanol/25% ammonia (100:10:1) as eluent, to give 285 mg of the title compound.

15 ¹H-NMR (CDCl₃, selected values): δ 0.55 (m, 1 H); 1.11, 1.12, and 1.17 (all s, together 6 H); 2.25 (s, 6 H); 2.45 (s, 3 H); 2.85 (s, 3 H); 4.02 (m, 1 H); 5.48 (dd, 1 H); 5.80 and 5.93 (m, and dd, together 1 H); 6.10 and 6.18 (both d, together 1 H); 6.87 and 7.00 (both m, together 1 H); 7.10 - 7.90 (m, 12 H).

20 HPLC 27.97 min (A1).
27.80 min (B1).


MS: 626.4 [M+1]⁺

25 For biological testing, the title compound was transferred into its diacetate salt by lyophilization from 0.5 acetic acid (40 ml).


Example 5

N-((1R)-1-{N-[(1R)-1-Benzyl-2-((2S)-2-((dimethylamino)methyl)pyrrolidin-1-yl)-2-oxoethyl]-N-methylcarbamoyl}-2-(2-naphthyl)ethyl)-N-methyl-3-((methylamino)methyl)benzamide

5



- 10 N-{3-[N-((1R)-1-{N-[(1R)-1-Benzyl-2-((2S)-2-((dimethylamino)methyl)pyrrolidin-1-yl)-2-oxoethyl]-N-methylcarbamoyl}-2-(2-naphthyl)ethyl)-N-methylcarbamoyl]benzyl}-N-methylcarbamic acid tert-butyl ester



15

At 0 °C, N-(3-dimethylaminopropyl)-N'-ethylcarbodiimide hydrochloride (132 mg, 0.69 mmol) was added to a solution of 3-(N-(tert-butoxycarbonyl)-N-methylamino)benzoic acid (183 mg,

0.69 mmol) and 1-hydroxy-7-azabenzotriazole (94 mg, 0.69 mmol) in dichloromethane (5 ml). The reaction mixture was stirred for 20 min at 0 °C. A solution of (2R)-N-[(1R)-1-benzyl-2-((2S)-2-((dimethylamino)methyl)pyrrolidin-1-yl)-2-oxoethyl]-N-methyl-2-(methylamino)-3-(2-naphthyl)propionamide (345 mg, 0.69 mmol) in dichloromethane (10 ml) and N,N-dimethylformamide (5 ml) and ethyldiisopropylamine (0.118 ml) were added successively. The reaction mixture was stirred for 16 h, while it was warming up to room temperature. It was diluted with ethyl acetate and washed with a saturated aqueous solution of sodium hydrogen carbonate (70 ml). The aqueous phase was extracted with ethyl acetate (3 x 50 ml). The combined organic layers were dried over magnesium sulfate. The solvent was removed in vacuo. The crude product was purified by flash chromatography on silica (40 g), using dichloromethane/methanol/25% aqueous ammonia (100:10:1) as eluent, to give 524 mg of N-{3-[N-((1R)-1-{N-[(1R)-1-benzyl-2-((2S)-2-((dimethylamino)methyl)pyrrolidin-1-yl)-2-oxoethyl]-N-methylcarbamoyl}-2-(2-naphthyl)ethyl)-N-methylcarbamoyl]benzyl}-N-methylcarbamamic acid tert-butyl ester.

¹H-NMR (CDCl₃, selected values): δ 0.72 (m, 1 H); 1.45 (br, 9 H); 3.18 (br, 6 H); 4.05 (m, 1 H); 4.32 and 4.40 (both br, together 2 H); 5.60 (dd, 1 H); 5.95 (m, 1 H); 6.80 - 6.90 (m, 16 H).

At 0 °C, trifluoroacetic acid (7 ml) was added to a solution of N-{3-[N-((1R)-1-{N-[(1R)-1-benzyl-2-((2S)-2-((dimethylamino)methyl)pyrrolidin-1-yl)-2-oxoethyl]-N-methylcarbamoyl}-2-(2-naphthyl)ethyl)-N-methylcarbamoyl]benzyl}-N-methylcarbamamic acid tert-butyl ester (523 mg, 0.70 mmol) in dichloromethane (7 ml). The reaction mixture was stirred for 25 min at 0 °C. The solvent was removed in vacuo. The residue was dissolved in dichloromethane (90 ml), and the solvent was removed in vacuo. The latter procedure was repeated two times. The crude product was purified by flash chromatography on silica (40 g), using dichloromethane/methanol/25% aqueous ammonia (100:10:1) as eluent, to give 436 mg of the title compound.

¹H-NMR (CDCl₃, selected values): δ 0.87 (m, 1 H); 1.22 (m, 1 H); 1.45 (m, 1 H); 1.67 (m, 1 H); 4.09 (m, 1 H); 5.53 and 5.90 (dd and m, together 2 H); 6.80 - 7.90 (m, 16 H).

HPLC 28.43 min (A1).
30.63 min (B1).

MS: 648.4 [M+1]⁺


5

For biological testing, the title compound was transferred into its diacetate salt by lyophilization from 0.5 M acetic acid (40 ml).

Example 6

10

(2E)-5-Amino-5-methylhex-2-enoic acid N-((1R)-1-{N-[(1R)-1-benzyl-2-(4-(dimethylamino)piperidin-1-yl)-2-oxoethyl]-N-methylcarbamoyl}-2-(2-naphthyl)ethyl)-N-methylamide.



15

The title compound was prepared as in example 1 using 4-(dimethylamino)piperidine hydrochlorid salt, (2R)-2-(N-(tert-butoxycarbonyl)-N-methylamino)-3-phenylpropionic acid, (2R)-2-(N-(tert-butoxycarbonyl)-N-methylamino-3-(2-naphthyl))propionic acid, and (2E)-5-(butoxycarbonylamino)-5-methylhex-2-enoic acid.


20

¹H-NMR : (CDCl₃; selected values) δ 1.40 (s, 6 H); 2.00 (s, 6 H); 4.42 - 4.85 (2 H); 5.45 - 5.90 (m, 2 H); 6.28 (dd, 1 H); 6.85(m, 1 H); 7.10 - 7.85(m, 12 H)

MS(ES) : m/z 626.2 (M+H)⁺

Example 7

- 5 (2E)-5-Amino-5-methylhex-2-enoic acid N-methyl-N-[(1R)-1-(N-methyl-N-[(1R)-1-[N-methyl-N-(1-methylpiperidin-4-yl)carbamoyl]-2-phenylethyl]carbamoyl)-2-(2-naphthyl)ethyl]amide




- 10 The title compound was prepared as in example 1 using 1-methyl-4-(methylamino)piperidine, (2R)-2-(N-(tert-butoxycarbonyl)-N-methylamino)-3-phenylpropionic acid, (2R)-2-(N-(tert-butoxycarbonyl)-N-methylamino-3-(2-naphthyl))propionic acid, and (2E)-5-(butoxycarbonylamino)-5-methylhex-2-enoic acid.
- 15 ¹H-NMR (CDCl₃; selected values) δ 5.50 - 6.08 (m, 2 H); 6.20 - 6.70 (m, 2 H); 7.10 - 7.85 (m, 12 H)

Example 8

3-Aminomethyl-N-((1R)-1-{N-[(1R)-1-benzyl-2-(4-methylpiperazin-1-yl)-2-oxoethyl]-N-methylcarbamoyl}-2-(2-naphthyl)ethyl)-N-methylbenzamide

5



The title compound was prepared as in example 1 using N-methylpiperazine, (2R)-2-(N-(tert-butoxycarbonyl)-N-methylamino)-3-phenylpropionic acid, (2R)-2-(N-(tert-butoxycarbonyl)-N-methylamino-3-(2-naphthyl))propionic acid, and 3-((tert-butoxycarbonylamino)methyl)benzoic acid.

¹H-NMR (CDCl₃; selected values) δ 3.30 (m, 1 H); 3.50 (dd, 1 H); 3.75 (m, 1 H); 3.95 (s, 2 H); 5.78 (t, 1 H); 3.88 (m, 1 H); 7.00 - 7.80 (16 H).

HPLC : 24.55 min (A1).

26.52 min (B1).


MS(ES) : m/z = 606.4 [M+H]⁺.

20

Example 9

(2E)-5-Amino-5-methylhex-2-enoic acid N-((1R)-1-{N-[(1R)-1-benzyl-2-(4-methylpiperazin-1-yl)-2-oxoethyl]-N-methylcarbamoyl}-2-(2-naphthyl)ethyl)-N-methylamide

5



The title compound was prepared as in example 1 using N-methylpiperazine, (2R)-2-(N-(tert-butoxycarbonyl)-N-methylamino)-3-phenylpropionic acid, (2R)-2-(N-(tert-butoxycarbonyl)-N-methylamino-3-(2-naphthyl))propionic acid, and (2E)-5-(butoxycarbonylamino)-5-methylhex-2-enoic acid.

¹H-NMR (CDCl₃; selected values) δ 1.24 (s, 6 H); 1.65 (s, 3 H); 2.35 (s, 3 H); 2.80 (s, 3 H); 5.68 (dd, 1 H); 5.78 (dd, 1 H); 6.18 (dd, 1 H); 6.95(m, 1 H); 7.15 - 7.80 (m, 12 H).


HPLC : 25.03 min (A1).
27.50 min (B1).

MS(ES) : m/z = 598.4 [M+H]⁺.

20

Example 10

(2E)-5-Amino-5-methylhex-2-enoic acid N-methyl-N-((1R)-1-{N-methyl-N-[(1R)-2-phenyl-1-((2,2,6,6-tetramethylpiperidin-4-yl)carbamoyl)ethyl]carbamoyl}-2-(2-naphthyl)ethyl)amide



5

The title compound was prepared as in example 1 using 4-amino-2,2,6,6-tetramethylpiperidin, (2R)-2-(N-(tert-butoxycarbonyl)-N-methylamino)-3-phenylpropionic acid, (2R)-2-(N-(tert-butoxycarbonyl)-N-methylamino-3-(2-naphthyl))propionic acid, and (2E)-5-(butoxycarbonylamino)-5-methylhex-2-enoic acid.

¹H-NMR (CDCl₃; selected values) δ 1.25 (s, 6 H); 1.40 (two s, 6 H); 1.52 (two s, 6 H); 2.92 (s, 3 H); 3.02 (two s, 3 H); 5.10 (dd, 1 H); 5.50 (dd, 1 H); 6.15 (d, 1 H); 6.75 (m, 1 H); 7.00 - 8.00 (m, 12 H).

HPLC : 29.27min (A1).


31.67min (B1).

MS(ES) : m/z = 654.8 [M+H]⁺.

20

Example 11

3-Aminomethyl-N-methyl-N-((1R)1-{N-methyl-N-[(1R)-2-phenyl-1-((2,2,6,6-tetramethylpiperidin-4-yl)carbamoyl)ethyl]carbamoyl}-2-(2-naphthyl)ethyl)benzamide



5

The title compound was prepared as in example 1 using 4-amino-2,2,6,6-tetramethylpiperidin, (2R)-2-(N-(tert-butoxycarbonyl)-N-methylamino)-3-phenylpropionic acid, (2R)-2-(N-(tert-butoxycarbonyl)-N-methylamino-3-(2-naphthyl))propionic acid, and 3-((tert-butoxycarbonylamino)methyl)benzoic acid.

10

¹H-NMR (CDCl₃; selected values) δ 3.60 - 3.85 (m, 2 H); 3.90 - 4.30 (m, 1 H); 5.25 - 5.95 (m, 2 H); 6.70 - 7.90 (m, 16 H).

15


HPLC : 29.27 min (A1).

31.55 min (B1).

MS(ES) : m/z = 662.4 [M+H]⁺.

Example 12

(2E)-5-Amino-3,5-dimethylhex-2-enoic acid N-methyl-N-((1R)-1-{N-methyl-N-[(1R)-2-phenyl-1-((2,2,6,6-tetramethylpiperidin-4-yl)carbamoyl)ethyl]carbamoyl}-2-(2-naphthyl)ethyl)amide



5

The title compound was prepared as in example 1 using 4-amino-2,2,6,6-tetramethylpiperidin, (2R)-2-(N-(tert-butoxycarbonyl)-N-methylamino)-3-phenylpropionic acid, (2R)-2-(N-(tert-butoxycarbonyl)-N-methylamino-3-(2-naphthyl))propionic acid, and (2E)-5-(butoxycarbonylamino)-3,5-methylhex-2-enoic acid.

10

¹H-NMR (CDCl₃; selected values) δ 3.92 - 4.30 (m, 1 H); 5.05 - 5.88 (m, 3H); 7.00 - 7.80 (m, 12 H).

15

HPLC : 29.80 min (A1).


32.43 min (B1).

MS(ES) : m/z = 668.4 [M+H]⁺.

20

Example 13

(2E)-4-(1-Aminocyclobutyl)but-2-enoic acid N-((1R)1-{N-[(1R)1-benzyl-2-(4-methylpiperazin-1-yl)-2-oxoethyl]-N-methylcarbamoyl}-2-(2-naphthyl)ethyl)-N-methylamide



5

The title compound was prepared as in example 1 using N-methylpiperazine, (2R)-2-(N-(tert-butoxycarbonyl)-N-methylamino)-3-phenylpropionic acid, (2R)-2-(N-(tert-butoxycarbonyl)-N-methylamino-3-(2-naphthyl))propionic acid, and (2E)-4-(1-(butoxycarbonylamino)cyclobutyl)but-2-enoic acid.

¹H-NMR (CDCl₃; selected peaks) δ 1.62 (s, 3 H); 2.35 (s, 3 H); 2.80 (s, 3 H); 5.70 (dd, 1 H); 5.80 (dd, 1 H); 6.22 (d, 1 H); 6.98 (m, 1 H); 7.15 - 7.80 (m, 12 H).

15

HPLC : 25.88 min (A1).
28.65 min (B1).


MS(ES) : m/z = 610.4 [M+H]⁺.

20

Example 14

(2E)-5-Amino-3,5-dimethylhex-2-enoic acid N-((1R)-1-{N-[(1R)-1-benzyl-2-(4-methylpiperazin-1-yl)-2-oxoethyl]-N-methylcarbamoyl}-2-(2-naphthyl)ethyl)-N-methylamide

5



- 10 The title compound was prepared as in example 1 using N-methylpiperazine, (2R)-2-(N-(tert-butoxycarbonyl)-N-methylamino)-3-phenylpropionic acid, (2R)-2-(N-(tert-butoxycarbonyl)-N-methylamino-3-(2-naphthyl))propionic acid, and (2E)-5-(butoxycarbonylamino)-3,5-methylhex-2-enoic acid.


- 15 ¹H-NMR (CDCl₃; selected values) δ 1.18 (s, 6 H); 1.68 (s, 3 H); 1.95 (s, 3 H); 2.30 (s, 3 H); 2.85 (s, 3 H); 3.40 (dd, 1 H); 3.54 - 3.75 (m, 2 H); 5.68 - 5.85 (m, 3 H); 7.15 - 7.80 (m, 12 H).

HPLC : 25.70 min (A1).
28.27 min (B1).

- 20 MS(ES) : m/z = 612.4 [M+H]⁺.

Example 15

(2E)-4-(1-Aminocyclobutyl)but-2-enoic acid N-((1R)-1-{N-[(1R)-1-benzyl-2-(4-hydroxypiperidin-1-yl)-2-oxoethyl]-N-methylcarbamoyl}-2-(biphenyl-4-yl)ethyl)-N-methylamide



This compound was prepared as in example 1 but using 4-hydroxypiperidine, (2R)-(N-tert-butoxycarbonyl-N-methylamino)-3-phenylpropionic acid and (2R)-2-(N-tert-butoxycarbonyl-N-methylamino)-3-(biphenyl-4-yl)propionic acid and (2E)-4-(1-(tert-butoxycarbonylamino)cyclobutyl)but-2-enoic acid as starting materials.

ESMS: 637.4 (M+H)⁺


HPLC: r_t = 33.58 min. (A1)

HPLC: r_t = 34.95 min. (B1)

Example 16

(2E)-5-Amino-5-methylhex-2-enoic acid N-((1R)-1-{N-[(1R)-1-benzyl-2-(4-hydroxypiperidin-1-yl)-2-oxoethyl]-N-methylcarbamoyl}-2-(biphenyl-4-yl)ethyl)-N-methylamide

5



This compound was prepared as in example 1 but using 4-hydroxypiperidine, (2R)-2-(N-tert-butoxycarbonyl-N-methylamino)-3-phenylpropionic acid and (2R)-2-(N-tert-butoxycarbonyl-N-methylamino)-3-(biphenyl-4-yl)propionic acid and (2E)-5-tert-Butoxycarbonylamino-5-methylhex-2-enoic acid as starting materials

10

ESMS: 625.4 (M+H)⁺

HPLC: r_t = 32.65 min. (A1)


15

HPLC: r_t = 34.02 min. (B1)

Example 17

(2E)-5-Amino-3,5-dimethylhex-2-enoic acid N-((1R)-1-{N-[(1R)-1-benzyl-2-(4-hydroxypiperidin-1-yl)-2-oxoethyl]-N-methylcarbamoyl}-2-(biphenyl-4-yl)ethyl)-N-methylamide

5



This compound was prepared as in example 1 but using 4-hydroxypiperidine, (2R)-2-(N-tert-butoxycarbonyl-N-methylamino)-3-phenylpropionic acid and (2R)-2-(N-tert-butoxycarbonyl-N-methylamino)-3-(biphenyl-4-yl)propionic acid and (2E)-5-tert-butoxycarbonylamino-3,5-dimethylhex-2-enoic acid as starting materials.

ESMS: 639.4 (M+H)⁺


HPLC: r_t = 33.29 min. (A1)

15 HPLC: r_t = 36.40 min. (B1)

Example 18

(2E)-5-Amino-5-methylhex-2-enoic acid N-((1R)-1-{N-[(1R)-1-benzyl-2-(4-hydroxypiperidin-1-yl)-2-oxoethyl]-N-methylcarbamoyl}-2-(2-naphthyl)ethyl)-N-methylamide

5



This compound was prepared as in example 1 but using 4-hydroxypiperidine, (2R)-2-(N-tert-butoxycarbonyl-N-methylamino)-3-phenylpropionic acid and (2R)-2-(N-tert-butoxycarbonyl-N-methylamino)-3-(2-naphthyl)propionic acid and (2E)-5-tert-Butoxycarbonylamino-5-methylhex-2-enoic acid as starting materials.

ESMS: 599.4 (M+H)⁺


HPLC: t_r = 29.88 min. (A1)

15

Example 19

(2E)-5-Amino-3,5-dimethylhex-2-enoic acid N-((1R)-1-{N-[(1R)-1-benzyl-2-(4-hydroxypiperidin-1-yl)-2-oxoethyl]-N-methylcarbamoyl}-2-(2-naphthyl)ethyl)-N-methylamide

5



This compound was prepared as in example 1 but using 4-hydroxypiperidine, (2R)-2-(N-tert-butoxycarbonyl-N-methylamino)-3-phenylpropionic acid and (2R)-2-(N-tert-butoxycarbonyl-N-methylamino)-3-(2-naphthyl)propionic acid and (2E)-5-tert-Butoxycarbonylamino-3,5-dimethylhex-2-enoic acid as starting materials.

10


ESMS: 613.4 (M+H)⁺

HPLC: r_t = 30.58 min. (A1)

Example 20

(2E)-4-(1-Aminocyclobutyl)but-2-enoic acid N-((1R)-1-{N-[(1R)-1-benzyl-2-(4-hydroxypiperidin-1-yl)-2-oxoethyl]-N-methylcarbamoyl}-2-(2-naphthyl)ethyl)-N-methylamide

5



This compound was prepared as in example 1 but using 4-hydroxypiperidine, (2R)-2-(N-tert-butoxycarbonyl-N-methylamino)-3-phenylpropionic acid and (2R)-2-(N-tert-butoxycarbonyl-N-methylamino)-3-(2-naphthyl)propionic acid and (2E)-4-(1-(tert-butoxycarbonylamino)cyclobutyl)but-2-enoic acid.

10


ESMS: 611.4 (M+H)⁺

HPLC: r_t = 30.82 min. (A1)

Example 21

(2E)-5-Amino-5-methylhex-2-enoic acid N-((1R)-1-{N-[(1R)-1-(4-fluorobenzyl)-2-(4-hydroxypiperidin-1-yl)-2-oxoethyl]-N-methylcarbamoyl}-2-(2-naphthyl)ethyl)-N-methylamide

5



This compound was prepared as in example 1 but using 4-hydroxypiperidine, (2R)-2-(N-tert-butoxycarbonyl-N-methylamino)-3-(4-fluorophenyl)propionic acid and (2R)-2-(N-tert-butoxycarbonyl-N-methylamino)-3-(2-naphthyl)propionic acid and (2E)-5-tert-butoxycarbonylamino-5-methylhex-2-enoic acid as starting materials.

ESMS: 617.4 (M+H)⁺


HPLC: $t_r = 30.27$ min. (A1)

15 HPLC: $r_t = 31.60$ min. (B1)

Example 22

(2E)-5-Amino-3,5-dimethylhex-2-enoic acid N-((1R)-1-{N-[(1R)-1-(4-fluorobenzyl)-2-(4-hydroxypiperidin-1-yl)-2-oxoethyl]-N-methylcarbamoyl}-2-(2-naphthyl)ethyl)-N-methylamide

5



This compound was prepared as in example 1 but using 4-hydroxypiperidine, (2R)-2-(N-tert-butoxycarbonyl-N-methylamino)-3-(4-fluorophenyl)propionic acid and (2R)-2-(N-tert-butoxycarbonyl-N-methylamino)-3-(2-naphthyl)propionic acid and (2E)-5-tert-butoxycarbonylamino-3,5-dimethylhex-2-enoic acid as starting materials.

ESMS: 631.4 (M+H)⁺


HPLC: r_t = 30.98 min. (A1)

15 HPLC: r_t = 32.38 min. (B1)

Example 23

(2E)-4-(1-Aminocyclobutyl)but-2-enoic acid N-((1R)-1-{N-[(1R)-1-benzyl-2-(4-(dimethyl-amino)piperidin-1-yl)-2-oxoethyl]-N-methylcarbamoyl}-2-(2-naphthyl)ethyl)-N-methylamide

5



The title compound was prepared as in example 1 using 4-N,N-dimethylpiperazine, (2R)-2-(N-(tert-butoxycarbonyl)-N-methylamino)-3-phenylpropionic acid, (2R)-2-(N-(tert-butoxycarbonyl)-N-methylamino)-3-(2-naphthyl)propionic acid, and (2E)-4-(1-(butoxycarbonyl-amino)cyclobutyl)but-2-enoic acid.

10

$^1\text{H-NMR}$ (CDCl_3 ; selected peaks) δ 1.90 (s, 3 H); 2.38 (s, 3 H); 2.45 and 2.47 (two s, 3 H) 2.78 and 2.80 (two s, 3 H); 6.32 (dd, 1 H); 6.90 (m, 1 H); 7.15 - 7.84 (m, 12 H).

15


HPLC : 26.72 min (A1).

MS(ES) : m/z = 638.4 $[\text{M}+\text{H}]^+$.

Example 24

(2E)-5-Amino-5-methylhex-2-enoic acid N-((1R)-1-{N-[(2R)-2-(4-hydroxypiperidin-1-yl)-2-oxo-1-((2-thienyl)methyl)ethyl]-N-methylcarbamoyl}-2-(2-naphthyl)ethyl)-N-methylamide

5



This compound was prepared as in example 1 but using 4-hydroxypiperidine, (2R)-2-(N-tert-butoxycarbonyl-N-methylamino)-3-(2-thienyl)propionic acid and (2R)-2-(N-tert-butoxycarbonyl-N-methylamino)-3-(2-naphthyl)propionic acid and (2E)-5-tert-butoxycarbonylamino-5-methylhex-2-enoic acid as starting materials.

10

ESMS: 605.4 (M+H)⁺


HPLC: r_t = 29.07 min. (A1)

15

Example 25

(2E)-5-Amino-3,5-dimethylhex-2-enoic acid N-((1R)-1-{N-[(2R)-2-(4-hydroxypiperidin-1-yl)-2-oxo-1-((2-thienyl)methyl)ethyl]-N-methylcarbamoyl}-2-(2-naphthyl)ethyl)-N-methylamide

5



This compound was prepared as in example 1 but using 4-hydroxypiperidine, (2R)-2-(N-tert-butoxycarbonyl-N-methylamino)-3-(2-thienyl)propionic acid and (2R)-2-(N-tert-butoxycarbonyl-N-methylamino)-3-(2-naphthyl)propionic acid and (2E)-5-tert-butoxycarbonylamino-3,5-dimethylhex-2-enoic acid as starting materials.


ESMS: 619.4 (M+H)⁺

HPLC: t_r = 29.76 min. (A1)

15

Example 26

(2E)-5-Amino-5-methylhex-2-enoic acid N-(((1R)-2-(biphenyl-4-yl)-1-{N-[(2R)-2-(4-hydroxypiperidin-1-yl)-2-oxo-1-((2-thienyl)methyl)ethyl]-N-methylcarbamoyl}ethyl)-N-methylamide




This compound was prepared as in example 1 but using 4-hydroxypiperidine, (2R)-2-(N-tert-butoxycarbonyl-N-methylamino)-3-(2-thienyl)propionic acid and (2R)-2-(N-tert-butoxycarbonyl-N-methylamino)-3-(biphenyl-4-yl)propionic acid and (2E)-5-tert-butoxycarbonyl-amino-5-methylhex-2-enoic acid as starting materials.

ESMS: 631.2 (M+H)⁺

HPLC: t_r = 32.20 min. (A1)

Example 27

(2E)-5-Amino-3,5-dimethylhex-2-enoic acid N-((1R)-2-(biphenyl-4-yl)-1-{N-[(1R)-2-(4-hydroxypiperidin-1-yl)-2-oxo-1-((2-thienyl)methyl)ethyl]-N-methylcarbamoyl}ethyl)-N-methylamide




This compound was prepared as in example 1 but using 4-hydroxypiperidine, (2R)-2-(N-tert-butoxycarbonyl-N-methylamino)-3-(2-thienyl)propionic acid and (2R)-2-(N-tert-butoxycarbonyl-N-methylamino)-3-(biphenyl-4-yl)propionic acid and (2E)-5-tert-butoxycarbonyl-amino-3,5-dimethylhex-2-enoic acid as starting materials.

ESMS: 465.4 (M+H)⁺

HPLC: t_r = 32.89 min. (A1)

Example 28

(2E)-5-Methyl-5-(methylamino)hex-2-enoic acid N-((1R)-1-{N-[(1R)-1-benzyl-2-(4-hydroxypiperidin-1-yl)-2-oxoethyl]-N-methylcarbamoyl}-2-(biphenyl-4-yl)ethyl)-N-methylamide



- 10 The title compound was prepared as in example 1 but using 4-hydroxypiperidine, (2R)-2-(N-tert-butoxycarbonyl-N-methylamino)-3-phenylpropionic acid and (2R)-2-(N-tert-butoxycarbonyl-N-methylamino)-3-(biphenyl-4-yl)propionic acid and (2E)-5-(N-(tert Butoxycarbonyl)-N-methylamino)-5-methylhex-2-enoic acid as starting materials


15

MS: m/z: 639.4 (M+H)⁺

HPLC: Method A1:R_t = 32.94 min

Example 29

(2E)-4-(1-Aminocyclobutyl)but-2-enoic acid ((1R)-1-{N-[(1R)-1-benzyl-2-(4-hydroxypiperidin-1-yl)-2-oxoethyl]-N-methylcarbamoyl}-2-(biphenyl-4-yl)ethyl)amide



5


HPLC: Rt = 31,55 min.(A1)

Rt = 33,11 min.(B1)

LC-MS: 623.6 [M+1]⁺

CLAIMS

1. A compound of the general formula I



formula I


10 wherein

R¹ is hydrogen or C₁₋₆-alkyl;


R² is hydrogen or C₁₋₆-alkyl;

15

L is



or



wherein R⁴ is hydrogen or C₁₋₆ alkyl;

20

p is 0 or 1;


q, s, t, u are independently from each other 0, 1, 2, 3 or 4;

r is 0 or 1;

the sum $q + r + s + t + u$ is 0, 1, 2, 3, or 4;

- 5 R^9 , R^{10} , R^{11} , and R^{12} are independently from each other hydrogen or C_{1-6} alkyl;

Q is $>N-R^{13}$ or



- 10 wherein o is 0, 1 or 2;


T is $-N(R^{15})(R^{16})$ or hydroxyl;

R^{13} , R^{15} , and R^{16} are independently from each other hydrogen or C_{1-6} alkyl;

15

R^{14} is hydrogen, aryl or hetaryl;

G is $-O-(CH_2)_k-R^{17}$,




20

wherein R^{17} , R^{18} , R^{19} , R^{20} and R^{21} independently from each other are hydrogen, halogen, aryl, hetaryl, C_{1-6} -alkyl or C_{1-6} -alkoxy;

k is 0, 1 or 2;

J is $-O-(CH_2)_l-R^{22}$,



- 5 wherein R^{22} , R^{23} , R^{24} , R^{25} and R^{26} independently from each other are hydrogen, halogen, aryl, hetaryl, C_{1-6} -alkyl or C_{1-6} -alkoxy;

l is 0, 1 or 2;

a is 0, 1, or 2;

10

b is 0, 1, or 2;

c is 0, 1, or 2;

15 d is 0 or 1;

e is 0, 1, 2, or 3;

f is 0 or 1;

20

R^5 is hydrogen or C_{1-6} -alkyl optionally substituted with one or more hydroxyl, aryl or hetaryl;

R^6 and R^7 are independently from each other hydrogen or C_{1-6} -alkyl, optionally substituted with one or more halogen, amino, hydroxyl, aryl, or hetaryl;

25

R^8 is hydrogen or C_{1-6} -alkyl, optionally substituted with one or more halogen, amino, hydroxyl, aryl, or hetaryl;

R^6 and R^7 or R^6 and R^8 or R^7 and R^8 may optionally form $-(CH_2)_i-U-(CH_2)_j-$, wherein i and j independently from each other are 1, 2 or 3 and U is $-O-$, $-S-$, or a valence bond;

5 M is arylene, hetarylene, $-O-$, $-S-$ or $-CR^{27}=CR^{28}-$;

R^{27} and R^{28} are independently from each other hydrogen or C_{1-6} -alkyl, optionally substituted with one or more aryl or hetaryl;


10 or a pharmaceutically acceptable salt thereof.

2. The compound according to claim 1 wherein R^1 is C_{1-6} -alkyl.

3. The compound according to any one of the preceding claims wherein R^2 is C_{1-6} -alkyl.

15

4. The compound according to any one of the preceding claims wherein L is



20 wherein R^4 is hydrogen or C_{1-6} alkyl;

p is 0 or 1;

q , s , t , u are independently from each other 0, 1, 2, 3 or 4;


25

r is 0 or 1;

the sum $q + r + s + t + u$ is 0, 1, 2, 3, or 4;

R^9 , R^{10} , R^{11} , and R^{12} are independently from each other hydrogen or C_{1-6} alkyl;

Q is $>N-R^{13}$ or




wherein o is 0, 1 or 2;

T is $-N(R^{15})(R^{16})$ or hydroxyl;

R^{13} , R^{15} , and R^{16} are independently from each other hydrogen or C_{1-6} alkyl; and

R^{14} is hydrogen, aryl or hetaryl.

5. The compound according to any one of the claims 1-3 wherein L is




wherein q, s, t, u are independently from each other 0, 1, 2, 3 or 4;

r is 0 or 1;

the sum $q + r + s + t + u$ is 0, 1, 2, 3, or 4;

R^9 , R^{10} , R^{11} , and R^{12} are independently from each other hydrogen or C_{1-6} alkyl;

Q is $>\text{N-R}^{13}$ or



5


wherein o is 0, 1 or 2;

T is $-\text{N}(\text{R}^{15})(\text{R}^{16})$ or hydroxyl;

10 R^{13} , R^{15} , and R^{16} are independently from each other hydrogen or C_{1-6} alkyl; and

R^{14} is hydrogen, aryl or hetaryl.


15 6. The compound according to any one of the preceding claims wherein G is



wherein R^{17} , R^{18} , R^{19} , R^{20} and R^{21} independently from each other are hydrogen, halogen, aryl, hetaryl, C_{1-6} -alkyl or C_{1-6} -alkoxy.


20

7. The compound according to any one of the preceding claims wherein J is




wherein R^{22} , R^{23} , R^{24} , R^{25} and R^{26} independently from each other are hydrogen, halogen, aryl, hetaryl, C_{1-6} -alkyl or C_{1-6} -alkoxy.

8. The compound according to any one of the preceding claims wherein M is arylene or
 5 $-CR^{27}=CR^{28}-$, wherein R^{27} and R^{28} independently from each other hydrogen or C_{1-6} -alkyl, optionally substituted with aryl or hetaryl.
9. The compound according to any one of the preceding claims wherein R^6 and R^7
 10 independently from each other are hydrogen or C_{1-6} -alkyl.
10. The compound according to any one of the claims 1-8 wherein R^6 and R^7 form $-(CH_2)_i-U-(CH_2)_j-$, wherein i and j independently from each other are 1, 2 or 3 and U is $-O-$, $-S-$, or a valence bond.
11. The compound according to any one of the preceding claims wherein R^8 is hydrogen or
 15 C_{1-6} -alkyl.
12. The compound according to any one of the preceding claims selected from
- 20 (2E)-5-Amino-5-methylhex-2-enoic acid N-((1R)-1-{N-[(1R)-1-benzyl-2-(4-((dimethylamino)methyl)piperidin-1-yl)-2-oxoethyl]-N-methylcarbamoyl}-2-(2-naphthyl)ethyl)-N-methylamide




(2E)-5-Amino-5-methylhex-2-enoic acid N-((1R)-1-{N-[(1R)-1-benzyl-2-((3S)-3-(dimethylaminomethyl)piperidin-1-yl)-2-oxoethyl]-N-methylcarbamoyl}-2-(2-naphthyl)ethyl)-N-methylamide



5


(2E)-4-(1-Aminocyclobutyl)but-2-enoic acid N-((1R)-1-{N-[(1R)-1-benzyl-2-((3S)-3-(dimethylaminomethyl)piperidin-1-yl)-2-oxoethyl]-N-methylcarbamoyl}-2-(2-naphthyl)ethyl)-N-methylamide



10


15

(2E)-5-Amino-5-methylhex-2-enoic acid N-((1R)-1-{N-[(1R)-1-benzyl-2-((2S)-2-((dimethylamino)methyl)pyrrolidin-1-yl)-2-oxoethyl]-N-methylcarbamoyl}-2-(2-naphthyl)ethyl)-N-methylamide



5


N-((1R)-1-{N-[(1R)-1-Benzyl-2-((2S)-2-((dimethylamino)methyl)pyrrolidin-1-yl)-2-oxoethyl]-N-methylcarbamoyl}-2-(2-naphthyl)ethyl)-N-methyl-3-((methylamino)methyl)benzamide




10

15


(2E)-5-Amino-5-methylhex-2-enoic acid N-(((1R)-1-{N-[(1R)-1-benzyl-2-(4-(dimethylamino)piperidin-1-yl)-2-oxoethyl]-N-methylcarbamoyl}-2-(2-naphthyl)ethyl)-N-methylamide.




5 (2E)-5-Amino-5-methylhex-2-enoic acid N-methyl-N-[(1R)-1-(N-methyl-N-[(1R)-1-[N-methyl-N-(1-methylpiperidin-4-yl)carbamoyl]-2-phenylethyl]carbamoyl]-2-(2-naphthyl)ethyl]amide




3-Aminomethyl-N-((1R)-1-{N-[(1R)-1-benzyl-2-(4-methylpiperazin-1-yl)-2-oxoethyl]-N-methylcarbamoyl}-2-(2-naphthyl)ethyl)-N-methylbenzamide




(2E)-5-Amino-5-methylhex-2-enoic acid N-((1R)-1-{N-[(1R)-1-benzyl-2-(4-methylpiperazin-1-yl)-2-oxoethyl]-N-methylcarbamoyl}-2-(2-naphthyl)ethyl)-N-methylamide




- 5 (2E)-5-Amino-5-methylhex-2-enoic acid N-methyl-N-((1R)-1-{N-methyl-N-[(1R)-2-phenyl-1-((2,2,6,6-tetramethylpiperidin-4-yl)carbamoyl)ethyl]carbamoyl}-2-(2-naphthyl)ethyl)amide




3-Aminomethyl-N-methyl-N-((1R)-1-{N-methyl-N-[(1R)-2-phenyl-1-((2,2,6,6-tetramethylpiperidin-4-yl)carbamoyl)ethyl]carbamoyl}-2-(2-naphthyl)ethyl)benzamide




(2E)-5-Amino-3,5-dimethylhex-2-enoic acid N-methyl-N-((1R)-1-{N-methyl-N-[(1R)-2-phenyl-1-((2,2,6,6-tetramethylpiperidin-4-yl)carbamoyl)ethyl]carbamoyl}-2-(2-naphthyl)ethyl)amide




(2E)-4-(1-Aminocyclobutyl)but-2-enoic acid N-((1R)-1-{N-[(1R)-1-benzyl-2-(4-methylpiperazin-1-yl)-2-oxoethyl]-N-methylcarbamoyl}-2-(2-naphthyl)ethyl)-N-methylamide



(2E)-5-Amino-3,5-dimethylhex-2-enoic acid N-((1R)-1-{N-[(1R)-1-benzyl-2-(4-methylpiperazin-1-yl)-2-oxoethyl]-N-methylcarbamoyl}-2-(2-naphthyl)ethyl)-N-methylamide




(2E)-4-(1-Aminocyclobutyl)but-2-enoic acid N-((1R)-1-{N-[(1R)-1-benzyl-2-(4-hydroxypiperidin-1-yl)-2-oxoethyl]-N-methylcarbamoyl}-2-(biphenyl-4-yl)ethyl)-N-methylamide



5


(2E)-5-Amino-5-methylhex-2-enoic acid N-((1R)-1-{N-[(1R)-1-benzyl-2-(4-hydroxypiperidin-1-yl)-2-oxoethyl]-N-methylcarbamoyl}-2-(biphenyl-4-yl)ethyl)-N-methylamide



10


123

(2E)-5-Amino-3,5-dimethylhex-2-enoic acid N-((1R)-1-{N-[(1R)-1-benzyl-2-(4-hydroxypiperidin-1-yl)-2-oxoethyl]-N-methylcarbamoyl}-2-(biphenyl-4-yl)ethyl)-N-methylamide



5


(2E)-5-Amino-5-methylhex-2-enoic acid N-((1R)-1-{N-[(1R)-1-benzyl-2-(4-hydroxypiperidin-1-yl)-2-oxoethyl]-N-methylcarbamoyl}-2-(2-naphthyl)ethyl)-N-methylamide




10

15


(2E)-5-Amino-3,5-dimethylhex-2-enoic acid N-((1R)-1-{N-[(1R)-1-benzyl-2-(4-hydroxypiperidin-1-yl)-2-oxoethyl]-N-methylcarbamoyl}-2-(2-naphthyl)ethyl)-N-methylamide



(2E)-4-(1-Aminocyclobutyl)but-2-enoic acid N-((1R)-1-{N-[(1R)-1-benzyl-2-(4-hydroxypiperidin-1-yl)-2-oxoethyl]-N-methylcarbamoyl}-2-(2-naphthyl)ethyl)-N-methylamide




(2E)-5-Amino-5-methylhex-2-enoic acid N-((1R)-1-{N-[(1R)-1-(4-fluorobenzyl)-2-(4-hydroxypiperidin-1-yl)-2-oxoethyl]-N-methylcarbamoyl}-2-(2-naphthyl)ethyl)-N-methylamide




125


(2E)-5-Amino-3,5-dimethylhex-2-enoic acid N-((1R)-1-{N-[(1R)-1-(4-fluorobenzyl)-2-(4-hydroxypiperidin-1-yl)-2-oxoethyl]-N-methylcarbamoyl}-2-(2-naphthyl)ethyl)-N-methylamide




5 (2E)-5-Amino-5-methylhex-2-enoic acid N-((1R)-1-{N-[(1R)-1-benzyl-2-(4-hydroxy-4-(2-thienyl)piperidin-1-yl)-2-oxoethyl]-N-methylcarbamoyl}-2-(2-naphthyl)ethyl)-N-methylamide



(2E)-5-Amino-5-methylhex-2-enoic acid N-((1R)-1-{N-[(1R)-1-(3-hydroxycyclohexylcarbamoyl)-2-phenylethyl]-N-methylcarbamoyl}-2-(2-naphthyl)ethyl)-N-methylamide




(2E)4-(1-Aminocyclobutyl)but-2-enoic acid N-((1R)-1-{N-[(1R)-1-benzyl-2-(4-(dimethylamino)piperidin-1-yl)-2-oxoethyl]-N-methylcarbamoyl}-2-(2-naphthyl)ethyl)-N-methylamide




5

(2E)-5-Amino-5-methylhex-2-enoic acid N-((1R)-1-{N-[(2R)-2-(4-hydroxypiperidin-1-yl)-2-oxo-1-((2-thienyl)methyl)ethyl]-N-methylcarbamoyl}-2-(2-naphthyl)ethyl)-N-methylamide




10 (2E)-5-Amino-3,5-dimethylhex-2-enoic acid N-((1R)-1-{N-[(2R)-2-(4-hydroxypiperidin-1-yl)-2-oxo-1-((2-thienyl)methyl)ethyl]-N-methylcarbamoyl}-2-(2-naphthyl)ethyl)-N-methylamide




127

(2E)-5-Amino-5-methylhex-2-enoic acid N-((1R)-2-(biphenyl-4-yl)-1-{N-[(2R)-2-(4-hydroxypiperidin-1-yl)-2-oxo-1-((2-thienyl)methyl)ethyl]-N-methylcarbamoyl}ethyl)-N-methylamide



5


(2E)-5-Amino-3,5-dimethylhex-2-enoic acid N-((1R)-2-(biphenyl-4-yl)-1-{N-[(1R)-2-(4-hydroxypiperidin-1-yl)-2-oxo-1-((2-thienyl)methyl)ethyl]-N-methylcarbamoyl}ethyl)-N-methylamide



10


15

(2E)-5-Methyl-5-(methylamino)hex-2-enoic acid N-((1R)-1-{N-[(1R)-1-benzyl-2-(4-hydroxypiperidin-1-yl)-2-oxoethyl]-N-methylcarbamoyl}-2-(biphenyl-4-yl)ethyl)-N-methylamide



5

(2E)-4-(1-Aminocyclobutyl)but-2-enoic acid ((1R)-1-{N-[(1R)-1-benzyl-2-(4-hydroxypiperidin-1-yl)-2-oxoethyl]-N-methylcarbamoyl}-2-(biphenyl-4-yl)ethyl)amide



10 and pharmaceutically acceptable salts thereof.

13. Use of a growth hormone secretagogue or a pharmaceutically acceptable salt thereof for the preparation of a medicament for the treatment of growth retardation in connection with asthma.

15

14. The use according to claim 13 wherein the growth hormone secretagogue is a compound according to any one of the preceding compound claims or a pharmaceutically acceptable salt thereof.

15. A pharmaceutical composition comprising, as an active ingredient, a compound according to any one of the preceding compound claims or a pharmaceutically acceptable salt thereof together with a pharmaceutically acceptable carrier or diluent.

5

16. A method of stimulating the release of growth hormone from the pituitary of a mammal, the method comprising administering to said mammal an effective amount of a compound according to any one of the preceding compound claims or a pharmaceutically acceptable salt thereof or of a composition according to any one of the preceding composition claims.

10

17. Use of a compound according to any one of the preceding compound claims or a pharmaceutically acceptable salt thereof for the preparation of a medicament for stimulating the release of growth hormone from the pituitary of a mammal.

15 18. Use of a growth hormone secretagogue or a pharmaceutically acceptable salt thereof for the preparation of a medicament for the treatment of growth retardation in connection with juvenile rheumatic arthritis or cystic fibrosis.

20 19. The use according to claims 13 or 18 wherein the growth hormone secretagogue is selected among growth hormone releasing peptides, growth hormone releasing peptidomimetics, and growth hormone releasing compounds of a nonpeptidyl nature.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/DK 99/00368

A. CLASSIFICATION OF SUBJECT MATTER

IPC6: C07K 14/60, C07K 5/02, C07K 5/06, A61K 38/05, A61K 38/25
According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC6: C07K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

SE,DK,FI,NO classes as above

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

REG, CAPLUS, WPI

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	WO 9858950 A1 (NOVO NORDISK A/S), 30 December 1998 (30.12.98) --	1-15,17-19
X	WO 9723508 A1 (NOVO NORDISK A/S), 3 July 1997 (03.07.97) --	1-15,17-19
A	WO 9803473 A1 (NOVO NORDISK A/S), 29 January 1998 (29.01.98) -- -----	1-15,17-19

☐ Further documents are listed in the continuation of Box C.

☒ See patent family annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"I" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

21 October 1999

Date of mailing of the international search report

22-10-1999

Name and mailing address of the ISA/

Swedish Patent Office

Box 5055, S-102 42 STOCKHOLM

Facsimile No. +46 8 666 02 86

Authorized officer

Carolina Gómez Lagerlöf/ELY

Telephone No. +46 8 782 25 00

INTERNATIONAL SEARCH REPORTInternational application No.
DK99/00368**Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)**

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 16
because they relate to subject matter not required to be searched by this Authority, namely:
See extra sheet.
2. ☒ Claims Nos.: 1-11 in part
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
The formulation of claims 1-11 is so complicated because of the long lists of cascading substituents that it does not comply with Article 6 PCT prescribing that claims shall be clear and concise. For this reason the search has mainly been limited to the examples.

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International application No.
DK99/00368

Claim 16 relate to a method of treatment of the human or animal body by surgery or by therapy/a diagnostic method practised on the human or animal body/Rule.39.1(iv). Nevertheless, a search has been executed for this claim. The search has been based on the alleged effects of the compound/composition.

INTERNATIONAL SEARCH REPORT

Information on patent family members

28/09/99

International application No.

PCT/DK 99/00368

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9858950 A1	30/12/98	AU 7906998 A	04/01/99
WO 9723508 A1	03/07/97	AU 1092997 A	17/07/97
		CA 2239711 A	03/07/97
		CN 1211991 A	24/03/99
		CZ 9801950 A	14/10/98
		EP 0869974 A	14/10/98
		HU 9802580 A	01/02/99
		IL 124702 D	00/00/00
		JP 11209336 A	03/08/99
		JP 11501054 T	26/01/99
		NO 982872 A	21/08/98
		PL 327227 A	07/12/98
WO 9803473 A1	29/01/98	AU 3434697 A	10/02/98
		EP 0923539 A	23/06/99
		US 5922770 A	13/07/99